

## Field trip 2

### Late Paleozoic and Mesozoic terrestrial environments in the Dolomites and surrounding areas

Evelyn Kustatscher<sup>1,2</sup>, Massimo Bernardi<sup>3,4</sup>, Fabio Massimo Petti<sup>3,5</sup>,  
Marco Avanzini<sup>3</sup> & Riccardo Tomasoni<sup>3</sup>

- <sup>1</sup> Naturmuseum Südtirol, Bindergasse 1, 39100 Bozen/Bolzano, Italy; e-mail: evelyn.kustatscher@naturmuseum.it;
- <sup>2</sup> Department für Geo- und Umweltwissenschaften, Paläontologie und Geobiologie, Ludwig-Maximilians-Universität and Bayerische Staatssammlung für Paläontologie und Geologie, Richard-Wagner-Straße 10, 80333 München, Germany;
- <sup>3</sup> MuseodelleScienze di Trento, Corsodel Lavoro edella Scienza 3, 38123 Trento, Italy; e-mail: massimo.bernardi@muse.it, fabio.petti@muse.it, marco.avanzini@muse.it, riccardo.tomasoni@muse.it;
- <sup>4</sup> School of Earth Sciences, University of Bristol, Bristol BS81RJ, UK;
- <sup>5</sup> PaleoFactory, Dipartimento di Scienze della Terra, Sapienza Università di Roma, P.le Aldo Moro 5, 00185, Roma, Italy.

#### 1 Topics and highlights of the excursion

The Southern Alps represent one of the classical areas for the study of Late Paleozoic to Early Mesozoic stratigraphy. Spectacular outcrops allow exceptional insights into the genesis of the Permian continental-marine succession, Middle–Upper Triassic carbonate platforms and Jurassic shallow-marine to coastal successions. This area played an important role in the history of geology: in the 18<sup>th</sup> century Giovanni Arduino introduced in the Venetian and Recoaro Alps the subdivisions that now identify the four Erathems (Primary to Quaternary). Déodat de Dolomieu collected in the Etschtal/Valle dell’Adige the carbonate rock called in 1792 *Dolomie* by Nicolas de Saussure (Zenger et al., 1994; McKenzie & Vasconcelos, 2009). The granitoid intrusions that metamorphosed the Middle Triassic limestone rocks in the area of Predazzo and Passo San Pellegrino played a major role in the demise, in the early 19<sup>th</sup> century, of the Neptunist Theory. Furthermore, since the late 19<sup>th</sup> century the Southern Alps and the Northern Calcareous Alps have been used to define the global Permian–Triassic stratigraphy and biostratigraphy (e.g., Richtigshofen, 1860;

Mojsisovics, 1879, 1882; Mojsisovics et al., 1895; Bittner, 1892; Brack et al., 2005, Mietto & Manfrin, 1995; Mietto et al., 2003, 2012). Since 1970, the number of studies in petrology, stratigraphy, sedimentology and palaeontology conducted in the area has increased substantially (Gianolla et al., 2009–2010; Roghi et al., 2014, and references therein) and the integration of numerous well-correlated sections has resulted in a reliable biostratigraphic framework (ammonoids, bivalves, conodonts, sporomorphs) tied to radioisotope age data and magnetic reversals (especially for the PT-boundary, and the Anisian–Ladinian and the Ladinian–Carnian boundary intervals; see Roghi et al., 2014 and references therein). At the same time, several studies based on discoveries made in the area allowed deriving new palaeoecological and palaeobiological models, which gave new insight of broad evolutionary interest (e.g. Schmidt et al., 2006; Posenato et al., 2014; Bernardi et al., 2015). Last but not least, since 2009 the Dolomites are included in UNESCO natural world heritage list which explicitly acknowledges the “spectacular landscapes, which reflect the complex geological architecture of the region” (Gianolla et al., 2009).

Although famous for its Triassic marine basinal and platform successions, the Dolomites include also important non-marine successions. The Athesian Volcanic Group represents the largest and best outcropping lower Permian volcanic area in Europe. Volcanic rocks are locally intercalated with sedimentary successions marking periods of volcanic inactivity. This alternation between radiometrically dated volcanic rocks and fluvio-lacustrine sedimentary successions gives the latter an exceptional time constrain. The early Permian Athesian Volcanic Group is covered by the late Permian fluvial Gröden/Val Gardena Sandstones. During the Triassic non-marine successions are rare. Terrestrial sediments can be found locally in the coastal Early Triassic Werfen Formation, the middle Anisian Piz da Peres Formation and Richthofen Conglomerate, subaerial settings of the middle Ladinian magmatic event and the middle-late Carnian paralic Travenanzes Formation. During the Early Jurassic the typical shallow-water limestones of the Calcari Grigi Group locally developed coastal facies that record terrestrial biota.

This excursion will take us to some of the most interesting outcrops of non-marine environments in the Dolomites and surrounding areas. This includes the late Permian Gröden/Val Gardena Sandstone of the Bletterbach gorge, the Middle and Late Triassic successions of the Etschtal/Valle dell'Adige and Nonstal/Val di Non (Bad Gfrill/Bagni di Caprile, Gampenpass/Passo Palade, Mt. Roen) and the Lower Jurassic successions of the famous Lavini di Marco dinosaur tracksite, near Rovereto. These sections, most of which intensively restudied in the last years by the authors of this guide, yielded well preserved plant remains, diverse tetrapod footprints as well as insect traces giving important insights into the paleoenvironmental and climatic conditions at the paleotropics during the late Palaeozoic and Mesozoic.

This field trip guide has been compiled by the authors based on personal data and observations but is largely based on several key publications of the same authors plus co-authors, namely: Avanzini et al. (1997, 2002a, 2013), Gennaro (2007), Petti et al. (2013), Roghi et al. (2014).

## 2 Geological and environmental settings

The stratigraphic framework of the Dolomites includes Permian to Cretaceous formations (for a more detailed overview see Gianolla et al., 2009), although the area is mostly famous for its Triassic successions, making the region a classical study area for the stratigraphy of this period (see Roghi et al., 2014). The stratigraphic basement is composed of lower Palaeozoic rocks, deformed and metamorphosed by the Carboniferous Variscan Orogeny (Brixen Quarzphyllite), covered by the first erosive event, the Waidbruck Conglomerate (Avanzini et al., 2007, 2012, 2013). An early Permian (trans-)tensional rifting resulted in an important volcanic activity that accumulated massive volcanic successions (Athesian Volcanic Group); during the periods of quiescence fluvio-lacustrine sediments deposited in small, local basins (Avanzini et al., 2007, 2013; Morelli et al., 2007; Marchetti et al., 2015). The middle Permian is missing due to an important unconformity, while the late Permian is mostly represented by fluvial red beds of the Gröden/Val Gardena Sandstone (Massari & Neri, 1997). A marine transgression from the East covered the area and triggered the accumulation of shallow marine evaporites and carbonates (Bellerophon Formation). The end-Permian mass extinction and its aftermath (Benton & Twitchett, 2003) are represented by the storm-dominated, shallow marine carbonate and terrigenous deposits of the basalmost Werfen Formation (e.g., Farabegoli et al., 2007). The strong tectonic activity of the Anisian basins (especially to the east) and emerged areas (especially to the west) (Bosellini, 1968), subjected the area to several emersion episodes and a differential subsidence (De Zanche et al., 1992, 1993; Gianolla et al., 1998a). These sequences are marked by subaerial unconformities, and formed by continental conglomerates, shallow-water terrigenous-carbonate deposits and finally deep-water sediments or prograding carbonate platforms. During the early Ladinian, the subsidence slowed down and the isolated nuclei expanded to large, up to thousand meters thick, platforms (Schlern/Sciliar Dolomite, Bosellini, 1984) while in the basins tens of metres of nodular-cherty limestones formed (Buchenstein/Livinallongo Formation). During the middle

Ladinian a violent massive intrusive and effusive magmatic activity in the Predazzo and Monzoni area formed locally subaerial areas, huge heterogeneous megabreccia bodies (Caotico Eterogeneo), pillow lavas and hyaloclastites. After the magmatic activity, the carbonate production restarted with a widespread progradation (Cassian Dolomite) and a high basinal sedimentation rate (St. Cassian/San Cassiano Formation). Later, the basins were filled by mixed terrigenous-carbonate deposits (Heiligkreuz Formation) that register several moist phases and yielded some of the oldest known amber with inclusions of microorganisms (e.g., Gianolla et al., 1998b; Schmidt et al., 2006). The return to aridity is registered in the middle-late Carnian by alternations of continental, paralic and shallow-marine mixed sediments (Travenanzes Formation). The decrease in siliciclastic input lead to the formation of a carbonate peritidal platform of regional extension (Hauptdolomit/Dolomia Principale; Bosellini & Hardie, 1988). In the Early Jurassic, shallow water limestones formed (Calcarei Grigi Group; Masetti & Bottoni, 1978; Sauro et al., 1995; Avanzini et al., 2007), while soon after the region drowned due to the passive margin evolution of the Adria microplate giving origin to pelagic successions with encrinites and condensed nodular limestones (Rosso Ammonitico; Masetti & Bottoni, 1978). Finally, during the Cretaceous, the deposition of deep-water clay, marls and micrites (Marne del Puez) was interrupted by the onset of turbiditic sand input (Flysch di Ra Stua), recording the early stages of the Alpine deformation (Sauro et al., 1995).

### 3 Overview of Excursion route and stops

This three-day fieldtrip starts from the Bletterbach gorge (Bozen/Bolzano Province), crosses the area of Nonstal/Val di Non and Etschtal/Valle dell'Adige and ends in the Lavini di Marco dinosaur tracksite (Rovereto, Trento Province) (Fig. 1). The fieldtrip illustrates the early Permian to Early Jurassic tectono-stratigraphic evolution of a distinctive sector of the Southern Alps. The first day mainly concerns the Permian period. In the

Bletterbach Gorge we will follow the evolution of this sector of the Alps from the uppermost portion of the Athesian Volcanic Group (Auer/Ora Formation; early Permian) to the Gröden/Val Gardena Sandstone (late Permian) fluvial deposits. During the second day we will move up from the Permian formations (Lana neighbourhood) to the Middle Triassic of the Gampenpass/Passo Palade observing a nearly complete stratigraphic succession. During several stops on the road from Lana to the pass (S.S. 238) we will see the different members of the Induan-Olenekian marine Werfen Formation, the carbonate and terrigenous Anisian units (Lower Sarl/Serla Dolomite, Piz da Peres Conglomerate, Voltago Conglomerate and Richthofen Conglomerate), the following upward mixed carbonate/terrigenous units (Gracilis Formation, Giovo Formation) and the late Illyrian carbonate platforms deposits (Contrin Formation). In the afternoon we will climb up Mt. Roen and, nearby Rifugio Oltradige (Hütte), we will stop by a spectacular outcrop of the Carnian Travenanzes Formation. From here we will also have a dramatic view of the western and central Dolomites that will be the focus of a session dedicated to landscape/panoramic geology. The last day is devoted to the Lower Jurassic of the Calcarei Grigi Group and particularly to the Hettangian-late Sinemurian units (Monte Zugna Formation and basal portion of the Rotzo Formation). We will have a trip to the Lavini di Marco site, exploring its geology and popular dinosaur trackways, and have the chance to talk about historical and present day research on the fossil plants of the Rotzo Formation.

#### **Day 1: The late Permian (Lopingian) flora and fauna of the Bletterbach**

The Bletterbach Gorge is located southeast of Bozen/Bolzano and is, since 2009, part of the Dolomites UNESCO. The beauty of the landscape, the colors of the rocks – especially on a late summer or autumn evenings – are exquisite and have given the gorge the nickname of “Gran Canyon of South Tyrol”. The Bletterbach is easily visible through walking paths that are used from spring to autumn (estimated 60.000 visitors each year); events and guided tours are coordinated by the visitor center of the Geoparc Bletterbach. The



Fig. 1: Excursion route of the field trip with the main stops of the three days.

succession exposed in the gorge starts at the bottom with the Cisuralian Athesian Volcanic Group and finishes at the top with the early Middle Triassic Contrin Formation preserving changing landscapes, climate and environmental conditions as well as the evolution of life, both plant and animals, through time.

#### Stop 1, The Auer/Ora Formation of the Athesian Volcanic Group

At the bottom of the Bletterbach Gorge is exposed the upper part of the Auer/Ora Formation (Fig. 2), the youngest volcanic unit of the Athesian Volcanic Group, earlier termed "Bozner Quarzporphyr". The top of the Auer/Ora Formation, dated about 274 mya, represents pyroclastic flow deposits accumulated in a tectonically controlled graben-like structure between Bozen/Bolzano and Auer/Ora (Morelli et al., 2007). The formation is composed of coherent and homogeneous welded

rhyolitic lapilli-tuffs of pink-red to orange-red colours. Up to 4 mm large sanidine, pink plagioclase and quartz phenocrysts float in a felsic matrix. Due to smaller phenocrysts and crystal fragments this groundmass appears heterogenous (Morelli et al., 2007). Fiamme formed of dark aphanitic or juvenile porphyric inclusions occur frequently. A regular network of subvertical joints and 8–10 spaced discontinuity lines cutting the rock parallel to the orientation of the fiamme are noticeable features at outcrop scale. The latter may be attributed to different ignimbrite flow units (Morelli et al., 2007).

#### Stop 2, The boundary between the Auer/Ora Formation and the Gröden/Val Gardena Sandstone

The base of the Gröden/Val Gardena Sandstone follows older Permian rhyolitic ignimbrites of the Auer/Ora Formation (Morelli et al., 2007). This boundary is associated to an unconformity of



Fig. 2: The lower sector of the Bletterbach showing the narrow valley with the steep slopes incised in the rhyolitic ignimbrites of the Auer/Ora Formation of the Athesian Volcanic Group.

about 14–27 Ma (Cassinis et al., 1999). The onset of the second Permian tectono-sedimentary cycle *sensu* Cassinis et al. (2002) initiated with filling up of a palaeorelief. The contact rhyolite-Gröden/Val Gardena Sandstone is exposed along the main gorge between the descent trail from the visitor centre and the waterfall (Fig. 3). By optic levelling of the top rhyolite between the main gorge and its tributaries, an upfilled palaeorelief of about 30 m can be reconstructed.

### Stop 3, Mature gypsisols in the lowermost part of the Gröden/Val Gardena Sandstone

The lowermost part of the Gröden/Val Gardena Sandstone represents a ~30 m thick succession of

exclusively terrestrial strata composed of stacked up to 8 m thick fining upwards cycles (Massari et al., 1994). Internally, these cycles start with up to 3 m thick coarse-grained sandstones or conglomerates at their bases and grade into shaly sands and silts towards their tops. Conglomerates, matrix supported conglomerates to conglomeratic sandstones, commonly form the base of the lower cycles whereas coarse-grained sandstones form the bases of the upper cycles. Conglomeratic beds often appear massive and unstratified and sandstones are made of horizontally laminated, low-angle cross-bedded and trough-cross bedded sandstone lithofacies. Both lithologies form tabular bodies that have a sharp but almost non-erosive base and show lateral shifts to finer



Fig. 3: Boundary (unconformity) between the Auer/Ora Formation and the Gröden/Val Gardena Sandstone.

grained lithofacies towards the West and South. Internal bounding surfaces are horizontally oriented or low-angle inclined; the resulting architectures of sandstone units appear sheet-like to splay-like, respectively. Comparable architectures are described from proximal parts of aggradational floodplains that underwent crevassing and sheet flooding. The splay-like sandstone bodies with low-angle inclined subordinated bounding surfaces resemble architectures of crevasse splays (for more details see Roghi et al., 2014) and the tabular and sheet-like sandstone bodies may originate from subaerial unconfined flows (Fisher et al., 2007).

Conglomerates and sandstones are followed by reddish to sometimes variegated shaly siltstones to silty sandstones. These sediments representing the upper part of individual cycles experienced substantial modification by pedogenic processes. Destratification due to repeated shrinking and swelling is very common and resulted in typical slickensides. However, the most prominent visible features are up to 3 m thick petrogypsic and petrocalcic horizons of mature gypsisols and calcisols. At the base of petrogypsic/petrocalcic horizons a few gypsum/carbonate nodules occur scattered, become larger and more common towards the top and finally coalesce in forming massive gypsum or carbonate crusts, often with an undulating top. The gypsisol exposed at the

floor of the Bletterbach is well recognised for its pseudo-anticlinal structures (gilgai relief) highlighted by secondary gypsum veins demonstrating that precipitation of gypsum followed vertic processes (Fig. 4). Thicknesses and maturities of gypsisols clearly show that crevassing and sheet flooding on the floodplains were followed by longer times of subaerial exposure under climates characterised by precipitation < evaporation.

#### Stop 4, The first marine incursion into the Gröden/Val Gardena Sandstone

The Gröden/Val Gardena Sandstone exposed in the valley cliffs below the Cephalopod Bank represents the transgressive part of sequence Lo 2 (*sensu* Posenato, 2000). In terms of cyclicities, lithofacies and sedimentary architecture the succession corresponds more or less to the succession exposed below. Up to 8 m thick fining upward cycles show successively decreasing thicknesses towards the Cephalopod Bank accompanied by a reduced number of pedogenic modified horizons. Sharp based sheet-like to splay-like sand bodies, dominantly formed of upper flow regime bed forms, grade vertically into shaly lithologies that comprise mature gypsisols, calcisols or vertisols, respectively. Accordingly, these tabular sand bodies are considered proximal crevasse splays and sheet sands of an aggradational floodplain. In contrast to that, a northern tributary of the




Fig. 4: Example of outcrop with mature gypsisols (from Roghi et al., 2014). (A) Gypsisol with few and scattered nodules at the base. To the top gypsum nodules become common, coalesce and form a massive petrogypsic horizon. (B) Prominent gypsisol showing pseudo-anticlinal structures (gilgai relief) highlighted by secondary gypsum veins. The lower part comprises gypsum nodules of a petrogypsic horizon. (from Roghi et al., 2014)

main gorge exposes an up to 7 m thick sandstone that is formed of sheet-like sigmoidal sand bodies that are separated by 15–30° inclined bounding surfaces. This indicates a lateral shifting architecture similar to the “epsilon” cross-bedding *sensu* Allen (1963) or shows a lateral-accretion macroform (LA) *sensu* Miall (1985). The lateral accreting sand body is interpreted as point bar sheets that formed due to subsequent migration of a high-sinuuous meandering channel. As both, the channel fill and floodplain deposits, are formed of bed load and suspended load the river type has been interpreted as mixed-load meandering river (Massari et al., 1994; Massari & Neri, 1997).

The short-term ingression evidenced by the Cephalopod Bank shifted this channel-floodplain system towards the West, and the Bletterbach area was transformed into a coastal-deltaic area. At the waterfall a ~3 m thick intercalation of thin silty sandstones and dark grey to black clays is exposed below the Cephalopod Bank (Fig. 5). The succession is bioturbated and rich in plant remnants. The 2.5 m thick Cephalopod Bank shows coarsening upwards from fine-grained to medium-grained sandstones. In the lower part, bedding appears horizontal with individual layers of wave ripples. Towards the top hummocky cross-bedding indicates successive shallowing (Massari et al., 1994). Above the Cephalopod Bank a plant rich intercalation of silty sandstones and dark clays, comparable to the intercalation below, is exposed. In the succession pedogenic features are lacking and lithofacies, bedding features and the high content of plant remains (the so called “cuticle horizon” of Clement-Westerhof, 1984, 1986, 1987; Poort & Kerp, 1990) suggest deposition under subaquatic conditions. The vertical succession of lithofacies can be best explained within a fluvio-deltaic setting. The succession around the waterfall is considered to represent the transition from a floodplain to an interdistributary bay and to a delta plain back to floodplain (for more details see Roghi et al., 2014).

#### Stop 5, Plant fossils around the Cephalopod bank


The first mention of plant fossils from the Bletterbach Gorge was by Perwanger (1946), who cited a “*Lepidodendron*-Stengel”. Later,

Leonardi (1948, 1968) figured “*Lepidodendron* cf. *sternbergi* Lindley et Hutton, *Schizolepis permensis* Heer, *Lepidodendron* cf. *veltheinianum* Sternberg, *Lepidodendron* sp., *Lebachia* (= *Walchia* auct.) *laxifolia* Florin and *Lebachia* (?) sp.” (Leonardi, 1948); most fragments were badly preserved. Researchers from Utrecht University started a detailed study in the area finding horizons yielding plant fragments with exceptionally preserved cuticles attributed to several conifer and seed fern taxa: *Ortiseia leonardii* Florin, 1964, *O. jonkeri* Clement-Westerhof, 1984, *O. visscheri* Clement-Westerhof, 1984, *Dolomitia cittertia* Clement-Westerhof, 1987, *Majonica alpina* Clement-Westerhof, 1987, *Pseudovoltzia liebeana* (Geinitz) Florin, 1927, *P. sjerpii* Clement-Westerhof, 1987 and *Peltaspermum martinsii* (Germar) Poort et Kerp, 1990. Bulk macerations revealed a flora dominated by conifers with seed ferns and ginkgophytes as additional elements (Kerp & Kustatscher, 2005) with a high percentage of gymnosperms, related to taphonomic selection. Several hundreds of better preserved plant remains were collected between 2003 and 2011 during a project of the Museum of Nature South Tyrol and the Geoparc Bletterbach. The plant fossils were assigned to 33 distinct fossil taxa, including foliage, stem fragments, fructifications, and dispersed seeds of sphenophytes, seed ferns, taeniopterids, ginkgophytes and conifers (Fig. 6) (Kustatscher et al., 2012, 2014a; Bauer et al., 2014; Roghi et al., 2014). Ginkgophyte leaf fragments are the dominant element (51%) including *Baiera digitata* (Brongniart) Heer, 1876, *Sphenobaiera* sp., putative *O-ha-tsuki*-type fertile leaves, seeds, and several leaves of uncertain affinities (Fischer et al., 2010; Bauer et al., 2014). The conifers represent the second most common plant group (~40% of the flora) with *Ortiseia leonardii* Florin, 1964, *Ortiseia visscheri* Clement-Westerhof, 1984, *Quadrocladus* sp. and *Pseudovoltzia* sp. The seed fern are rare in the flora (*Sphenopteris suessii* Geinitz, 1869, *Sphenopteris* sp., *Germaropteris martinsii* (Germar in Kurtze, 1839) Kustatscher et al., 2014 (Poort & Kerp, 1990; Kustatscher et al., 2014a) as are taeniopterid leaf fragments (< 1.5%) and horsetails (0.2%). The plant remains show also indications of plant-animal interactions (Roghi et al., 2014), although with a low frequency of insect damage patterns.

Fig. 5:  
The Cephalopod Bank in the Bletterbach Gorge.



Fig. 6:  
(a) Leaf of *Baiera digitata*;  
(b) Leaf fragment of *Taeniopteris* sp. B.  
(c) Shoot fragment of *Ortiseia leonardii*;



## Stop 6, The tetrapod footprints of the Bletterbach Gorge

The first report of a fossil fauna in the Bletterbach Gorge area dates back to the 19th century when Ernst Kittl collected a single tetrapod footprint downstream the gorge, nearby the village of Egna (Kittl, 1881; see also Abel, 1929). The first deliberate palaeontological explorations were organized in the 1940s by Piero Leonardi who surveyed the canyon looking for plants and footprints (e.g., Leonardi 1951, 1953, 1968). Excavations continued in the following years and numerous tetrapod tracks were collected from distinct fossiliferous horizons (e.g., Conti et al., 1975, 1977, 1980; Ceoloni et al., 1988; Leonardi & Nicosia, 1973; Leonardi et al., 1975; Wopfner, 1999). From then on the Bletterbach became globally known for its Loapingian tetrapod association (just to cite a few international fieldtrips run in this locality see Conti et al., 1986, Pittau et al., 2005; Gianolla et al., 2010; Roghi et al., 2014).

Thirteen footprint-bearing horizons were recognized within 180 m thick succession corresponding to the Lo 1–3 sequences of Posenato (2010). Most of these findings come from the lower and middle part of the Gröden/Val Gardena Sandstone (first 120 meters), but some footprint-rich beds have been found within the continental deposits interfingering with the marine sediments of the overlying Bellerophon Formation, at the top of the 3rd depositional sequence recognized in the upper Permian succession. One of the most productive horizons lays some hundred meters upstream the waterfall. To date, only footprints document the presence of tetrapod vertebrates, since no skeletal remains have been found. The ichnoassemblage is however abundant and shows a high diversity allowing a detailed reconstruction of the vertebrate fauna. The abundance of fossil footprints has probably been favoured by their high preservation potential, due to the sedimentological context and high sedimentation rate (Avanzini & Tomasoni, 2004).




Fig. 7: The Gröden/Val Gardena Sandstone trophic network as derived from the study of vertebrate ichnoassociation, fossil plants and plant-arthropod interaction (from Roghi et al., 2014)

The ichnofauna counts thirteen ichnotaxa belonging to various groups of synapsid (therapsids), parareptiles (pareiasaurs), eureptiles (captorhinids and neodiapsids as younginiformes), and archosauriformes (see Conti et al., 2000; Avanzini et al., 2011; Bernardi et al., 2015). The Bletterbach ecosystem was characterised by large-sized primary consumers (pareiasaurs, diadectomorpha) that possibly fed on high-fibrous plants, such as ginkgophytes and conifers that constitute the largest part of the floral association. Small herbivores (captorhinids, small therapsids) were probably effective in shredding and crushing plant material. Large sized carnivorous predators (archosauriformes, gorgonopsids) seem to be less abundant, even though a preservational bias cannot be excluded. Small secondary consumers (undetermined neodiapsids) were probably carnivorous-insectivores and would have fed on the entomofauna documented by foliage insect

feeding traces described above (Fig. 7) (Bernardi et al., 2014).

Among the various features of the vertebrate association we here highlight two key components, which give particular relevance to this locality worldwide. The site contains (1) the (abundant) presence of pareiasaur footprints, and (2) the occurrence of the oldest archosaur footprints worldwide. One of the most common fossil footprints in the Bletterbach association is *Pachypes dolomiticus* Leonardi et al., 1975, a track attributed to pareiasaurs (Leonardi et al., 1975; Valentini et al., 2008, 2009). *Pachypes* is the only known footprint convincingly referred to pareiasaurs (Valentini et al. 2009). About 40 specimens (tracks and trackways) are known to date from this site, all preserved as natural casts (Fig. 8). The other exceptional features of the Gröden/Val Gardena Sandstone outcropping in this area are




Fig. 8: Two key features of the Bletterbach ichnofauna. a, c: The well preserved pareiasaur tracks *Pachypes dolomiticus* Leonardi et al., 1975, abundant in the Gröden/Val Gardena Sandstone succession. b, d: Two chirotheriid footprints attributed to archosaurian trackmakers, from Leonardi et al. (1975); Conti et al. (1977); Valentini et al. (2009); Bernardi et al. (2015).

chirotheriid footprints (Conti et al., 1977; Wopfner, 1999; Bernardi et al., 2015). These tracks, assigned to “Chirotheriid indet.” or to cf. *Protochirotherium*, can be confidently attributed to archosauriform trackmakers (Bernardi et al., 2015). Archosauriforms radiated from the Lower Triassic, but the occurrence of chirotheriid tracks in the upper Permian (Lopingian) of the Dolomites indicates that derived archosauriforms were already present before the PT-boundary. Together with the occurrence of the recently discovered fragmentary archosauriform *Eorasaurus* from the upper Permian of Russia (Ezcurra et al., 2014), these tracks provided evidence for a broader evolutionary radiation of archosauriforms in the late Permian than expected (Bernardi et al., 2014, 2015), further highlighting the scientific relevance of the Bletterbach palaeontological site worldwide.

### **Day 2 morning: The Triassic of the Nonstal/Val di Non area**

The area crossed by the field trip on the second day is interested by a monoclinical, gently dipping towards Southwest (Fig. 9).

In this geographic sector volcanic and sedimentary units regularly overlap each other (Avanzini et al., 2001, 2007, 2012). An important tectonic lineament (Völlan/Foiana Line) borders to the west the Gallberg/Monte Gall-Shonegg/Macaion block bringing into contact the sedimentary units of the Etschtal/Valle dell’Adige with the volcanic Laugen Spitze/Monte Luco units (Fig. 10). Close to the Völlan/Foiana Line the monoclinical is characterised by a series of narrow folds with subvertical flanks and axis parallel or gently dipping (en-echelon)

with respect to the master fault. A low-angle fault set in the Middle Triassic clastic units cuts the summit portion of Gallberg/Monte Gall and is connected to the Völlan/Foiana Line along the Brandis creek. The Southern Alps succession of this sector includes units from the metamorphic basement of Variscan age up to the Middle Triassic sedimentary units (essentially marls and limestone). The Southern Alps basement, buried beneath the alluvial deposits of Adige River near Lana, is composed of quartz-phyllite intruded by lower Permian plutons and dykes (Monte Cross Granodiorite near Lana). The basement is covered by a thick lower Permian volcanic succession (1500–2000 m), recently mapped as Athesian Volcanic Group, coheval and comagmatic with the above mentioned intrusions. The Permo-Mesozoic sedimentary cover has a relatively reduced thickness in comparison to the Western Dolomites and Venetian Prealps. Furthermore, they are characterised by several gaps, particularly in the Triassic (upper Ladinian–Carnian) with peculiar sedimentological features. The area between the Etschtal/Valle dell’Adige and the Giudicarie, of which the Mendel/Mendola–Schöneegg/Macaion chain represents the northernmost sector, is interpreted as a relative structural high for the whole Mesozoic. The substrate is followed by a Pleistocene-Holocene cover articulated in a variety of diverse geological bodies mainly related to the glacial dynamics that during the Quaternary strongly controlled the geological evolution of this area. These deposits are well represented in the lateral valleys, where they establish complex heteropic and overlapping relationships (Avanzini et al., 2013).




Fig. 9: The Nonstal/Val di Non and the excursion route. It mainly follows Avanzini et al. (2013).

Stop 1, The San Ippolito hill and the lower Permian volcanic rocks

The hill is modelled within the Gargazzone Formation and shows clearly the effect of the extensive

glacial exaration related to the last Wurmian pleniglacial. The bump is constituted by a wide exposed surface of sheepback rocks. Volcanic rocks are represented by black to dark grey




Fig. 10: Geological section across the Völlan/Foiana Line near the Gampenpass/Passo Palade.



Fig. 11: The top of the Permian volcanites: The Gargazzone Formation at San Ippolito Hill.



Fig. 12: Outcrop at the base of the Gröden/Val Gardena Sandstone near Narano.

pyroclastic ignimbrite with red-to-violet tones along the cracks due to its progressive oxidation. The surfaces exposed to subaerial weathering show colors varying from light grey to orange. From a stratigraphical point of view we are at the top of the lower Permian volcanites (Fig. 11). The wide plateau that extends toward S–SE, including the village of Tisens/Tesimo, marks the top of the volcanic succession, cropping out in the lower escarpments that overlook the Etschtal/Valle dell'Adige. The boundary between the volcanites and the upper sequences is marked by a regional erosional unconformity; the clastic deposits of the Gröden/Val Gardena Sandstone therefore cover different stratigraphic units of the Athesian Volcanic Group.

#### Stop 2, The lower portion of the Gröden/Val Gardena Sandstone

Here the typical lithofacies of the lower Gröden/Val Gardena Sandstone are characterised by grey-to-white sandstones intercalated with levels of coarser sandstone and microconglomerates (Fig. 12). Fining upward sequences of medium-to-coarse sandstone, with low-angle cross lamination can be recognized. They are organised into thin lenticular units marked by a sharp erosional base. In the basal portion these units show conglomerate lenses made by centimetre-sized quartz clasts. The depositional environment of this portion of the Gröden/Val Gardena Sandstone can be referred to a proximal fluvial system constituted by a net of shallow channels characterised by low sinuosity.

### Stop 3, The top of the Gröden/Val Gardena Sandstone and the base of the Werfen Formation

The top of the Gröden/Val Gardena Sandstone, between the Etschtal/Valle dell'Adige and the Giudicarie, is characterised by red, grey and whitish sandstone alternating with red or grey siltites organised in sandy bars and channel-fill sandstone bodies (Fig. 13). The topmost 3–3.5 metres of the formation are made by an alternation of black sandstone and arenaceous dolostone with mm- to cm-sized interlayers of silty marls. Above the last sandy bank lies a succession of finely stratified reddish pelites (< 1m). Carbonate intraclasts and nodules ("caliche"), related to the occurrence of paleosols at the top of the unit, are clearly visible within this pelitic succession. A sandy-to-silty unit with ripple marks covers the top of the formation. This unit is 1 m thick and is constituted by a dense alternation of pelites and fine sandstone; the dominant sedimentary structures are




Fig. 13: The upper part of the Gröden/Val Gardena Sandstone.

current- and wave-generated ripples. Bioturbation is represented by vertical burrows. This shallow water unit is followed, through an erosional surface, by sandy deposits and finally by the oolitic layers of the basal Werfen Formation (Tesero Member; Induan *p.p.*). This interval replaces the Bellerophon Formation along the northwestern Etschtal/Valle dell'Adige (Cassinis et al., 1993).

### Stop 4, The base of the Werfen Formation

The base of the Werfen Formation (late Changhsingian–Olenekian *p.p.*) is represented by the Tesero Member (Induan *p.p.*), grey-yellowish to brown coarse-grained dolostone with a clear oolitic texture (Fig. 14). At the base, the banks are massive and up to 50–60 m thick, showing an internal subdivision in thinner layers welded by stilolitic joints. The depositional environment of the Tesero Member is shallow marine (subtidal) allowing the development of oolitic bodies and their reworking by the wave. Temporary changes in the hydraulic regime toward milder hydrodynamic conditions allowed the settling of the finer horizons.

Above the massive banks, the Tesero Member gradually changes into an alternation of grey-yellowish marly limestone and marls. The marly limestone, organised in cm-thick laminae or more rarely into dm-thick layers, contain a small quartzitic-micaceous silty fraction, often bioturbated and with nodular structures. These layers are intercalated with marly limestones and marls with graded bioclastic lenses. In both lithotypes parallel lamination is frequent, while low-angle cross lamination is rare. These layers correspond to the Mazzin Member that reflects a different depositional environment with muddy seabeds and low hydrodynamic energy.

### Stop 5, From the Andraz Horizon to the Campil Member (Werfen Formation)

In this stop the Andraz Horizon (Induan *p.p.*) is formed by a dense alternation of yellowish marly or marly-silty evaporitic dolostones and reddish siltstones. This horizon is approximately 7 m thick and completely devoid of fossil remains. The lower boundary is marked by sudden increase of the silty fraction. The dolostones are arenaceous



Fig. 14: The basal part of the Tesero Member within the Werfen Formation.

and granular. Vacuolar texture is common in all horizons and the surface beds are characterised by desiccation cracks (mud-cracks, small teepees). The depositional environment is reconstructed as an arid tidal flat with fluctuations of inter-supratidal conditions. This is the first peritidal episode of the Werfen Formation and marks the top of the first of four major sedimentary cycles with regressive trend documented within the formation itself. The overlying Seis/Siusi Member (Induan *p.p.*-Olenekian *p.p.*) is made of an alternation of grey and/or reddish marly and marly-silty limestones, sometimes bioturbated, with oolitic-bioclastic calcarenites ("*Gastropoden-Oolith*"). The calcarenites (grainstone and packstone) are made of oolites, bivalves and microgastropods; more rarely they are characterised by a high percentage of recrystallized peloids (packstone). The pelitic lithotypes constitute cm-sized laminae or

more rarely they form massive banks about 1 m thick while calcarenites layers are cm to dm-sized. An evaporitic horizon occurs at the top of the unit, composed of some metres thick reddish to yellowish silty dolostones and siltstones. They represent the upper boundary of the unit that ends the second sedimentary cycle with a regressive trend documented within the formation. The depositional environment is a shallow marine seabed, characterised by high wave and recurrent storm events that allowed the deposition of bioclastic sandstone.

Above is the Gastropods Oolith (Olenekian *p.p.*), made up of reddish oolitic-bioclastic calcareous sandstones rich in bivalves and small gastropods. Strata are few cm to 10 cm thick and are intercalated with gray marls and breccia horizons ("*Koken-conglomerat*"). Bioturbation is frequent. Total thickness of the member is 20–25 meters.




Fig. 15: Part of the Werfen Formation outcropping at the bridge at km. 24 of the S.S. 238.

The depositional environment is reconstructed as a shallow sea, above wave base. In the upper part of the succession a gradual transition to the Campil Member (Olenekian *p.p.*) crops out. It is marked by an alternation of decimetric red sandstones and reddish siltstones or silty marls. The latter are organised in cm-sized layers. The sedimentary structures encompass different types of cross- and parallel-lamination, and load structures (ball-and-pillows). The depositional environment is a muddy seabed, below the wave base level but always less deep than the underlying members (Fig. 15).

#### Stop 6, The Anisian successions

The Anisian succession is constituted by terrigenous and terrigenous-carbonate units essentially deposited in basinal, lagoonal, peritidal and continental environments. Three terrigenous, mainly conglomeratic, units of subsequent ages (Piz da Peres Conglomerate, Voltago Conglomerate and Richthofen Conglomerate, each representing the base of a third-order sequence) are followed upward by some mixed carbonate/terrigenous units (i.e. Gracilis Formation, Giovo Formation) and by carbonate platform deposits (Contrin Formation) (Figs. 16, 17).




Fig. 16: Stratigraphic section of the sedimentary units of Anisian age between Nonstal/Val di Non and Etschtal/Valle dell'Adige and their correlation with those of the northeastern Dolomites. SLI – Lower Sarl/Serla Dolomite; PPS, Piz da Peres Conglomerate; GLS – Gracilis Formation; VTG – Voltago Conglomerate; GIV1 – Gampenpass/Passo Palade Member of the Giovo Formation; GIV2, Kurtatsch/Cortaccia Member of the Giovo Formation; GIV3 – Monte Ozol Member of the Giovo Formation; REC – Recoaro Limestone; SLS – Upper Sarl/Serla Dolomite; RIC – Richthofen Conglomerate; MRB – Morbiac Limestone; CTR – Contrin Formation.

The Lower Sarl/Serla Dolomite (late Olenekian-early Anisian) is composed of stratified light-grey dolostones (10–30 cm thick) with stylolites. The dolostones are intercalated with marly horizons following the cyclicity in a peritidal environment, with inter-supertidal horizons marked by reddish surfaces. The depositional environment is a shallow tidal flat. The boundary with the overlying Gracilis Formation (Bitinian–early Pelsonian) is marked by an erosional surface (paraconformity); the Gracilis Formation is a carbonate-terigenous sequence made by intercalations of calcareous-marly dolostones and slightly nodular bioclastic calcareous sandstones. Its upper part (ca. 40 meters) is made of grey-yellow silty limestones. The boundary with the Lower Sarl/Serla Dolomite is sharp and easily recognisable, being marked by also by colour change from white

dolostones below to the grey lithofacies above. The Piz da Peres Conglomerate of the eastern Dolomites is missing. The Gracilis Formation documents a depositional environment with tidal flats and small carbonatic platforms near coastal environments with predominantly terrigenous sedimentation (Fig. 17).

The middle Anisian (Pelsonian) palaeogeographic and stratigraphic setting of the Southern Alps is complex. The sedimentary units show a great variability in thickness and facies associations (Fig. 18). Along the Etschtal/Valle dell'Adige, the Pelsonian units show a wide lateral continuity and thickness, the base of which is constituted by Voltago Conglomerate (?Bithynian–earliest Pelsonian), which can be up to 130 m thick and finally the Giovo Formation.





Fig. 17: Early–middle Anisian paleogeographic evolution of the Nonstal/Val di Non and Etschtal/Valle dell'Adige.

Fig. 18:  
Correlation scheme between the formations of the lower-middle Anisian and the sedimentary evolution of Nonstal/Val di Non and Etschtal/Valle dell'Adige.



The last Anisian carbonate platform is constituted by the Contrin Formation (late Illyrian). In the Etschtal/Valle dell'Adige and surrounding areas the Pelsonian sedimentary units are thick and laterally extended while the Illyrian terrigenous successions are represented only locally and by thin and immature conglomerates. The great carbonate platforms of the Dolomites are almost completely absent.

#### Stop 7, The Giovo Formation

The Giovo Formation is a composite formation of up to 140 m thickness, typical of the Etschtal/Valle dell'Adige. It can be distinguished in three members: Gampenpass/Passo Palade Member (GIV1), Kortatsch/Cortaccia Member (GIV2), Monte Ozol Member (GIV3).

The Gampenpass/Passo Palade Member (GIV1) represents, locally, the base of the formation. It is made of yellow to gray sandstones with carbonate cement and graded conglomerates with cross lamination. Fossil content, although

discontinuous, comprises abundant plant remains and crinoids (*Encrinus liliiformis*, *Dadocrinus gracilis*). The clasts, made of a variety of carbonates, evaporitic dolostones and clastic units derive from the top of the underlying Voltago Conglomerate (Fig. 19). Sandstone horizons can be alternated with grey pelites, often bioturbated and rich in plant remains. Cross lamination at different scale can be recognized together with slumpings. Thickness varies from 0 to 50 m. The depositional environment is marginal marine (foreshore) with high terrigenous input from the emergent land.

The Kortatsch/Cortaccia Member (GIV2) overlaps or locally replaces the Gampenpass/Passo Palade Member. In the lower portion it is composed of siltstones and grey marls that are rich in plant remains. Above there is an alternation of peritidal limestones and silty-sandstones, similar to the "Calcari scuri del Torrente Framont" (Agordo Formation) described by Pisa et al. (1979). Strata have variable thickness (5–20 cm) and are often heavily bioturbated, resulting in the general nodularity of



Fig. 19: Typical aspect to the Gampenpass/Passo Palade Member at Unsere Liebe Frau im Walde/Se-nale.

the interval. Above, subtidal limestones are more frequent. They can be represented by bioturbated (nodular) calcareous sandstones, rich in ostracods, crinoids and brachiopods, or by encrinets (e.g., Gallberg/Monte Gall Tunnel, along S.S. 238) (Fig. 20). Thickness of the member ranges from 20 to 50 m. The lowermost horizons are rich in plant remains, invertebrate traces (*Rhizocoralium*) and vertebrate tracks (*Rhynchosauroides*, *Synaptichnium*, *Chirotherium*). Calcareous algae (*Diplopora* sp.) and forams (*Glomospira* sp.) are often associated with ostracods, brachiopods (*Tetractinella trigonella*, *Rhynchonella decurtata*, *Coenothyis vulgaris*), corals (*Montlivaltia*, *Thamnasteria* cf. *silesiaca*) and crinoids (*Encrinus liliiformis*, *Dadocrinus gracilis*). The depositional environment can be related with a complex setting of lagoons and shallow seaways with frequent emersion phases that were gradually replaced laterally by an open sea with carbonate sedimentation typical of a subtidal platform.

Fig. 20: Kurtatsch/Cortaccia Member at the tunnel at km 15.9 of the S.S. 238.




Fig. 21: Dark dolomites at the base of the Monte Ozol Member near Gampenpass/Passo Palade.



Fig. 22: *Encrinurus liliiformis* fragments in the outcrop near the tunnel of km 15.9 of the S.S. 238.



Fig. 23: Carbonate lithofacies of the Gracilis Formation near the tunnel at km 15.9 of the S.S. 238.




Fig. 24: Scheme of the Anisian outcrops along the S.S. 238 near Sankt Felix/San Felice.

The Monte Ozol Member (GIV3) was considered the base of the Mendel-Dolomit by Richthofen (1874). It is made of dolostones, dark gray bioclastic limestones and calcareous sandstones. They are organized in 10 cm thick strata, moderately bioturbated and are replaced upward by subtidal grey dolostones (Fig. 21). Bioclasts are mostly crinoid fragments, forming locally entire horizons of encrinites. The thickness of the member varies from 10 to 40 m. The boundary with the overlying Contrin Formation can be marked by an erosional surface (e.g., Shonegg/Macaion) and few cm of volcanic clay. These strata can be correlated with the Richthofen Conglomerate that was not deposited in this area. This interval crops out along the road just before the tunnel, where a wide outcrop of white dolostones can be observed.

The Giovo Formation crops out near the tunnel, where a sub-vertical transpressive fault puts into contact the Kurtatsch/Cortaccia Member of the Giovo Formation (in which the tunnel was dug) with its underlying Gampenpass/Passo Palade Member (just above the tunnel). Outside the tunnel the Kurtatsch/Cortaccia Member can be

observed with its typical disarticulated crinoid remains (*Encrinus liliiformis*) (Fig. 22).

About 200 meters uphill a second sub-vertical fault borders a sequence of sub-vertical strata dipping eastward. At this outcrop the boundary between the Gracilis Formation (Fig. 23) and the underlying Lower Sarl/Serla Dolomite can be observed. This is the last carbonate unit in the area, before crossing the Völlan/Foiana fault after which the lower Permian volcanites of Laugen Spitze/Monte Luco crops out.

Stop 8, The Voltago conglomerate and the base of the Giovo Formation

The Voltago conglomerate crops out in the high Nonstal/Val di Non and in the Etschtal/Valle dell'Adige along the Mendelkamm/Mendola mountain chain. It was first described by Kreis (1971) and corresponds to the "Mittlere Pereschichten" of Bechstädt & Brandner (1971) (for more details see Avanzini et al., 2001, 2007). The Voltago Conglomerate (Pelsonian *p.p.*) is characterised by discontinuous conglomeratic horizons

of up to 2 m thickness. These intervals are chaotic near the base, while towards the top they show gradation and cross lamination. The lowermost portion is composed of white dolomitic clasts (Lower Sarl/Serla Dolomite) in a whitish-yellow matrix. Silty limestones and grey or reddish arenaceous dolostones follow. Upward follows a bioturbated red-grey sandstones that near the top is replaced by an intercalation of stromatolitic dolostones and arenaceous evaporitic dolostones alternated with green or red siltstones with a thickness of ca. 10 cm. Within these horizons thin smectites document the original deposition of volcanic ashes. Parallel (or low angle) laminations, ripples and mud cracks are the main sedimentary structures of the unit. Bioturbation is abundant. The conglomerate reaches up to 130 m in thickness. (Fig. 24)

This site yielded an abundant association of tetrapod tracks with *Parasynaptichnium gracilis*, *Synaptichnium pseudosuchoides*, *Isochirotherium delicatum* and *Rhyncosauroides tirolicus* (Avanzini, 2000, 2002b). Plant remains belong to conifer shoots of the genus *Voltzia*. The depositional environment is terrestrial to marginal marine. In the lower portion of the formation it documents the development of a deltaic system and vegetated swamps, which may evolve in lagoons. In the upper part the environment shifts to a carbonate tidal flat influenced by strong evaporation. The Voltago Conglomerate is separated from the Giovo Formation (Gampenpass/Passo Palade Member) by an evaporitic lithozone (Fig. 25). The lower portion of the Gampenpass/Passo Palade Member is here visible and represented

by cross-laminated yellow-grey sandstones with carbonate cement. Some micro-conglomerates show clasts of evaporitic dolostones which source is the top of the Voltago Conglomerate. The depositional environment is a marginal marine foreshore with a high terrigenous supply from the emergent lands.

#### Stop 9, The Anisian fauna of Nonstal/Val di Non - Roen

The Anisian formations of the Nonstal/Val di Non and the Mendelkamm/Mendola-Roen mountain chain yielded a well-preserved vertebrate ichno-association that documents the presence of a rich variety of reptile taxa (Avanzini & Neri, 1998; Avanzini, 2000, 2002b; Valdiserri & Avanzini, 2007) (Fig. 26). The footprints found in the Voltago Conglomerate of the northernmost sector of Nonstal/Val di Non and Tisens/Tesimo Valley (Gampenpass/Passo Palade, Bad Gfrill/Bagni di Caprile, Unsere liebe Frau in Walde/Senale) can be mostly referred to lepidosauromorph and archosaurian producers. The most abundant morphs can be assigned to the widespread ichnogenus *Rhyncosauroides* and most frequently to the species *R. tirolicus* (Avanzini & Renesto, 2002). These document the presence of lepidosauromorph reptiles from 10–60 cm in length. Less common, but in all well represented, are medium- to large-sized tracks assigned to the chirotheriid ichnogenes *Synaptichnium*, *Chirotherium*, cfr. *Brachychirotherium* and *Isochirotherium* (Avanzini & Lockley, 2002; Avanzini & Mietto, 2008a,b). These tracks can be attributed to archosaur producers and document the presence of whole communities of



Fig. 25: Change from red sandstones (on the left) to evaporitic dolomites (on the right) along the S.S. 238.

Fig. 26: Some of the tetrapod footprints of the Voltago Conglomerate: A - *Synaptichnium* sp. with impressions of the skin and *Brachychirotherium* sp. (Unsere Liebe Frau im Walde/Senale); B - *Isochirotherium delicatum* (Unsere Liebe Frau im Walde/Senale); C - *Chirotherium barthi* (Bad Gfrill/Bagni di Caprile); D - *Rotodactylus* sp. (bottom left) and *Rhyncosauroides tirolicus* (top right) (Bad Gfrill/Bagni di Caprile). Scale bar = 2 cm.


trakmakers, distributed along age classes. Reconstruction of the population structure for the trackmaker of *I. delicatum*, for example, was made possible by the discovery, in the Gampenpass/Passo Palade are of hundreds of slabs with exceptionally preserved fossil specimens (Avanzini & Lockley, 2002). Exceptionally, fine anatomical details of the tracks (e.g., skin/scales) were also preserved at times (Avanzini, 1999).

Just few kilometers southeast, in the deep Hölental/Val di Inferno that cuts the eastern flank of Monte Roen, near the village of Tramin/Termeno, other chirotheroid tracks have been described by Avanzini & Leonardi (2002). The unique morphological features allowed the erection of a new ichnospecies: *Isochirotherium infernii*. The track-bearing horizons, of Illirian age, were originally considered to be part of the Morbiac Limestone but are now attributed to the Giovo Formation (Petti et al., 2013). The abundance of these findings and the tight correlation between marine and continental associations allowed developing a tetrapod biochronology scheme for the Anisian in this area (Avanzini & Mietto, 2008a).

All these findings have special ecological interest given the paucity of skeletal remains of Anisian age discovered in the Dolomites are so far. With the exception of few isolated fragmentary remains (see Petti et al., 2013), the record of continental tetrapods is in fact only represented by a well-preserved lepidosauromorph, *Megachirella wachtleri* (Renesto & Posenato, 2003; Renesto & Bernardi, 2014).

#### Stop 10, The Anisian flora of Nonstal/Val di Non

The oldest flora appearing in the fossil record of the Triassic of the Dolomites shows a highly diversified vegetation dominated by ferns, cycadophytes and conifers. The floras of the Dolomites and Recoaro areas consist of typical Middle



Triassic elements such as the osmundaceous ferns (*Neuropteridium*, *Anomopteris*, *Gordonopteris*), herbaceous and subarborescent lycophytes (*Lepacyclotes*, *Lycopia*, *Isoetites*) and conifers (*Voltzia*, *Albertia*, *Pelourdea*; e.g., Broglio Loriga et al., 2002). Seedferns (*Scytophyllum*, *Peltaspernum*, *Sagenopteris*) and cycadophytes (*Bjuvia*, *Nilssonina*, *Dioonitocarpidium*) are common, while horsetails are rare (*Equisetites*; e.g., Broglio Loriga et al., 2002; Kustatscher et al., 2014a).

In the Nonstal/Val di Non the plant remains are represented by stem and strobili fragments of horsetails (*Equisetites conicus* Sternberg, 1833), frond fragments of ferns (*Neuropteridium elegans* (Brongniart) Schimper, 1879, *Gordonopteris lorigae* van Konijnenburg-van Cittert et al., 2006), reproductive organs of cycads (*Dioonitocarpidium*

sp. cf. *Dioonitocarpidium pennaeformis* (Schenk) Rühle von Lilienstern, 1928) and shoot fragments of conifers (*Voltzia walchiaeformis* Fliche, 1908, *Voltzia recubariensis* Schenk, 1868) (Fig. 27).

Although the plant fossils from the Nonstal/Val di Non are very fragmentary and not very well preserved, they however contain some of the typical Anisian elements from the Southern Alps such as the ferns *Gordonopteris lorigae*, *Neuropteridium elegans* and the conifers *Voltzia recubariensis* and *Voltzia walchiaeformis*. It is worth to mention the conifer *Voltzia recubariensis* occurrence, which a typical element of the Anisian floras of the whole Southern Alps (Brack & Kustatscher, 2013). The cycadophyte megasporophyll *Dioonitocarpidium* is also common in the Anisian of the Southern Alps while it appears in the German Basin only with the Ladinian (e.g., Kustatscher et al., 2013). *Equisetites conicus* was so far the first record from the Anisian; this taxon is generally known from the Ladinian of the German Basin and the Carnian of the Alpine area (Kustatscher et al., 2013).

### Day 2 afternoon: The Late Triassic of the Roen succession

The Mt. Roen outcrop is located west of the village of Tramin/Termenò (Bozen/Bolzano Province) in the uppermost Höllental/Val di Inferno, at an altitude of 2000 m a.s.l.

The outcrop can be referred to the Travenanzes Formation (=upper portion of the "Raibl Beds" Auct.; Neri et al., 2005). The Travenanzes Formation shows a strong lateral variation in this area, as typical of transitional and coastal environment with interfingering between terminal fan/flood plain and a shallow lagoonal environment with periodic input of continental detritic sediments (D'Orazi Porchetti et al., 2008) (Fig. 28).

Samples of black and grey shales from the lowermost part of the Travenanzes Formation have yielded a remarkable amount of organic matter, mainly composed of amorphous material and sporomorphs (Gennaro, 2007). According to Roghi (2004), the palynological association belongs to the *Granuloperculatipollis rudis* assemblage, characterised by Circumpolles *Partitisporites*




Fig. 27: Some constituents of the Anisian flora found in this area. a) *Neuropteridium* sp.; b) ovuliferous cone scale; c, d) *Voltzia recubariensis*. Schenk, 1868




Fig. 28: Palaeogeographic reconstruction of the region during the deposition of the Travenanzes Formation (from Gennaro, 2007).

*quadruplicis* and *Granuloperculatipollis rudis*. This latter species always occurs with the “long range” elements *Enzonalasporites vigens*, *Pseudoenzonalasporites summus*, *Camerosporites secatus* and *Duplicisporites verrucosus* (Roghi, 2004). Apart from *Partisporites quadruplicis* all other five species have been found in the collected samples, often in association with *Ricciisporites tuberculatus*. This association indicates a Tuvalian age.

#### Stop 1, The Travenanzes Formation at Monte Roen

In the Roen area the Travenanzes Formation mainly consists of interbedded white-gray aphanitic to silty dolostones and reddish or greenish shales, with subordinate sandstone to conglomerate intercalations (Fig. 29). This mixed siliciclastic-carbonate succession has been interpreted as representing a marginal marine environment with supplies of terrigenous or fully marine sediments, albeit shallow.

Facies alternations suggest interfingering between alluvial-plain, carbonate tidal flat and shallow-lagoon deposits (Avanzini et al., 2010, 2012; Breda & Preto, 2011). The upper portion of the Travenanzes is organized into meter-scale, and is constituted of shallowing-upward peritidal cycles, not easily distinguishable from those of the overlying Hauptdolomit/Dolomia Principale (Breda & Preto, 2011). In the classical central/eastern Dolomites sections the Travenanzes Formation lies on the Heiligkreuz Formation (lower Carnian) while in other areas as those crossed by this field trip (i.e., Nonstal/Val di Non, Etschtal/Valle dell’Adige) it lies directly on the Schlern/Sciliar Dolomite (upper Anisian–upper Ladinian) or on volcanics (upper Ladinian), with a sharp and erosional contact. In the Monte Roen sector the Travenanzes Formation is constituted by terrigenous, carbonate and evaporitic units deposited in fluvial, lagoonal and peritidal environments.




Fig. 29: The Monte Roen section (from Gennaro, 2007).

### Stop 2, The Travenanzes Formation ichnoassociation

The Travenanzes ichnoassociation is very diversified, and is dominated by medium-to-large archosaur tracks (*Brachychirotherium*, *Evazoum*, cf. *Atreipus*, *Grallator*) (Fig. 30). The related faunal association is well diversified including aetosaurs, crocodile-like archosaurs, dinosauromorphs and dinosaurs (D’Orazi Porchetti et al., 2008; Avanzini et al., 2010; Bernardi et al., 2010, 2013). In the Roen section two dinosaur footprints have been found about 6 m above the base of the Travenanzes Formation. The track-bearing layer lies within

an interval of 50 cm, characterised by whitish aphanitic dolostones separated by thin levels of red and green shales (Gennaro, 2007).

The tracks can be confidently attributed to theropod dinosaur trackmakers with an estimated body length of about 5 m, and a body mass value approximately of 200 kg. This size is not known from dinosaur skeletal remains of Carnian age found worldwide (Benton, 2006; Brusatte et al., 2010) and is approximated only by the Norian theropod dinosaur *Liliensternus* (Cuny & Galton, 1993; Lucas et al., 2006). These specimens are part of a diverse Carnian ichnofauna discovered in the area, as in




Fig. 30: A - Theropod pedal imprint from the Monte Roen site. B – the finding is confirmed by a similar specimen found in the Mostizzolo site, just few km south. Both tracks were found in the Travenanzes Formation (from Bernardi et al., 2013).

the Mostizzolo (near Cles village, about few km south west from Monte Roen; Bernardi et al., 2010) and San Gottardo outcrops (D’Orazi Porchetti et al., 2008). The Travenanzes faunal assemblage of Southern Alps has thus been considered one of the most diversified for the Tuvlian interval worldwide (Bernardi et al., 2013) and documents one of the earliest forms of evidence of dinosaurs in northern Pangea, and the co-occurrence with dinosauriforms and crurotarsal archosaurs provides support to the non-abrupt replacement of the archaic fauna by a dinosaurian one (Bernardi et al., 2013).

### Stop 3, The Travenanzes Formation flora

The Carnian flora is generally rich and diversified, a typical example being the Lunz flora in Austria. This flora experiences generally the first sure occurrence of bennettitalean together with the horsetails, ferns, cycads, seedferns and conifers. In the Travenanzes and San Cassian Formation plant remains are poorly preserved and generally restricted to horsetail stem fragments (e.g. *Equisetites*), fern frond fragments (e.g., *Danaeopsis*) and conifer shoot fragments. Recently discovered floras of Carnian age from the Julian Alps (Dogna) and the Dolomites (Rifugio Dibona)

yield some conifer shoots possibly attributed to the Cheirolepidiaceae while the cuticles of other conifer shoots resemble the Permian conifers (see Roghi et al., 2006). In general, the dominant group are the conifers (*Voltzia*, *Brachyphyllum*).

### **Day 3: The Early Jurassic: the Lavini di Marco ichnosite and the flora of the Rotzo Formation**

The Lavini di Marco tracksite is located few kilometers south of Rovereto (Trento Province) and represents one of the most important European dinosaur footprint-bearing outcrops. It has been studied thoroughly by several authors that, since 1990s, carried out extensive ichnological, sedimentological and palynological research (Lanzinger & Leonardi, 1992; Leonardi & Avanzini, 1994; Avanzini et al., 1997; Leonardi & Mietto, 2000; Avanzini et al., 2001, 2002a, 2003; Piubelli et al., 2005; Avanzini et al., 2006).

### **Stop 1, The Lavini landslide**

Few natural phenomena in the Alps have been the subject such controversial geologic, historic, philological and literary interpretations as the Lavini di Marco (Venzo, 2000). The descriptions of the Lavini are mostly confined to the interpretation of the causes of the event or to the debate on the probable mention of it in Dante's *Divina Comedia* (1314). The name Lavini or Slavini comes from the late Latin word *labina* which means "landslide" (Fig. 35).

Geological studies started in the XIX century and focused on attempts to date and identify the origin of the event. The accumulation was considered to be the result of multiple, different events and according to some authors, it would have destroyed the village of Lagaris or Lagare (Venzo, 2000). Orombelli & Sauro (1988) identify at least two events: The main one, which took place in 883 A.D., and another much older landslide, that



Fig. 31: The Lavini di Marco site and its landslide




Fig. 32: Palaeogeographic restoration of the central-eastern sector of Southern Alps during the Early Jurassic (From Petti et al., 2011c).

occurred more than 4000 years ago. The dinosaur footprints are preserved on the wide bed surfaces (“*laste*”) exposed by the falling of the overlying strata (Fig. 31).

#### Stop 2, Geology and palaeogeography: The Calcarei Grigi Group and the Trento Platform

The Lavini di Marco trampled horizons have been ascribed to the middle-upper part of the Monte Zugna Formation (Hettangian–Sinemurian *p.p.*) “Middle Peritidal Unit” *sensu* Avanzini et al. (2006) within the Calcarei Grigi Group (Bosellini & Loriga, 1971) and cover approximately 300.000 m<sup>2</sup> of monoclinical surfaces (Piubelli et al., 2005). From a palaeogeographic point of view, the Calcarei Grigi Group belong to the Trento carbonate Platform, now extending north–south from Verona to Bozen/Bolzano (NE Italy) and that during the Mesozoic covered an area of approximately 20.000 km<sup>2</sup> (Fig. 32).

This domain was characterised by shallow-water carbonate sedimentation through all the Early Jurassic and was bounded on the east by the Belluno pelagic basin and on the west by the Lombard pelagic basin through the so-called “Garda escarpment”, a normal fault system active during the Jurassic and the Cretaceous (Castellarin et al., 1993, 2005). As suggested by palynological analysis of some track-bearing horizons (Avanzini et al., 2006), the climate during the Early Jurassic varied from arid to humid conditions. The discovery in 1989 of the Lavini di Marco tracksite gave rise to an extensive search of other dinosaur tracks-bearing outcrops in the Calcarei Grigi Group. As a result to date twelve dinosaur tracksites were discovered to different formations of the Calcarei Grigi Group (Fig. 33) (see Avanzini & Petti, 2008 for a review).




Fig. 33: Lithostratigraphic successions of the Lower Jurassic Calcarei Grigi Group, showing the stratigraphic positions of the different dinosaur tracksites.

### Stop 3, Sedimentology of the track-bearing strata

The lowermost part of the Monte Zugna Fm. is made of alternations of subtidal limestones in metric beds and levels of green clay, without evidence of tidal deposits. By contrast, peritidal cycles with structures typical of the tidal environment characterise the upper part of the member where the dinosaur

footprints are found. The subtidal facies, representing the lower 2/3 of the cycle, are mainly made up of homogeneous micritic, peloidal-fossiliferous limestones, completely reworked by bioturbation. Sometimes accumulations of large bivalves are present. There are both storm layers and colonies with individuals in life position (*Gervillia buchii*).

The supratidal facies are characterised by peculiar sedimentary structures such as storm-bacterial lamination (stromatolites), mud cracks, teepees, birdseyes, flat pebble breccias etc. (Masetti, 2000). The fossils in this member are rare. Brachiopods at the base of the Monte Zugna Formation are represented by "*Terebratula*" *dubiosa* Haas and *Pisirhynchia uhligi* Haas. Small foraminifers (Textulariidae, Ataxophragmiidae, *Mancyna* cf. *termieri*) are found at various levels. The most important microfossil from a chronostratigraphical point of view is the alga *Palaeodasycladus mediterraneus* (Pia), which is an Early Jurassic marker. Together with the early Sinemurian ammonoid *Charmasseiceras* in the uppermost underlying Loppio Formation it constrains the age of the track-bearing layers to the Hettangian (Avanzini et al., 2006).

In the Lavini di Marco ichnosite, seven dinoturbaed levels have been recognized in a 7 meter-thick succession (Avanzini et al., 2002a). The richest footprint levels are numbered 104, 105 and 106. Layer 104 refers to the upper 10–15 cm of a 70 cm thick subtidal bed. It is a bioclastic packstone to rudstone, which grades upwards to a peloidal wackestone containing ostracods. Locally, the topmost surface shows small-scale depressions, from a few mm to 100 mm deep, separated by hillocks. The depressions are commonly filled by laminated, ostracod-bearing mudstone and wackestone, which are capped by dolomitic crust, clay, and iron oxides. The "ridges" consist of typical, subtidal bioclastic packstone and grainstone. These microtopographic features seem indicative of lateral karstification by analogy with observations on modern tropical carbonate islands (Borrouhil-Le Jean, 1993). The influence of freshwater is indicated by the stable isotope trend and, in particular, by  $\delta^{18}\text{O}$  values below -2‰ (Land, 1986). At the top of layer 104 there are few trackways exhibiting excellently preserved theropod footprints.

Layer 105 consists of alternating stromatolitic laminae and light gray peloidal mudstones, dark gray bioclastic wackestones and reddish mudstones. The lower boundary is marked by a light brown, laminated mudstone which smooths the irregular surface topography of layer 104. A continuous layer of dark-gray wackestone,

about 10 mm thick, containing *Thaumatoporella parvovesiculifera* is intercalated within the inter-supratidal stromatolitic bindstones. This continuous layer, probably a storm layer, grades into hazel-colored mudstone spotted by iron oxides, and then it is capped by a fine grained breccia or by a laminated mudstone-wackestone with evidences of pedogenetic rubefaction. This rubefacted horizon is characterised by iron-rich glaebules (Bain & Foos, 1993), clotted micrite and circumgranular cracking, all features indicative of supratidal conditions. SEM observations and EDAX analyses revealed that the red, pedogenetic horizons consist of a mixture of dolomite, limestone, clay and iron oxides. Smaller theropod footprints are common where layer 105 is thinner. Larger, deeper theropod and sauropod tracks, occur in thicker parts.

Layer 106 (100–120 mm thick; Fig. 34) exhibits lateral facies variations in the various sectors of the outcrop. The boundary with layer 105 is marked by dark gray stromatolitic bindstone and fenestral mudstone with iron oxides concentrated between cyanobacterial laminae. The top of the stromatolitic layer is pervasively dolomitized in the northern sectors of the outcrop. Study of the upper part of layer 106 allows the identification of a series of microenvironmental belts. At the Colatoio Chemini fine-grained, mud-supported breccias with disrupted laminated mudstone clasts, black pebbles, and light-brown subtidal lithoclasts crop out. Eastward, these breccias grade in to mudcracked mudstones which are locally affected by strong rubefaction. Further




Fig. 34: A polished section across layer 106, which preserves the majority of fossil footprints.

east and southeast, fenestral mudstones pass to coarser sediments which exhibit wave ripples. The mudcracked, rubefacted sediments limit the occurrence of pervasive dolomitization and establish a boundary between two environments characterised by different footprints. The dolomitized sector exhibits deep sauropod footprints and tracks that cut across the entire layer 106 and part of layer 105. Exceptionally, some footprints extend down to the top of layer 104.

#### Stop 4, The ichnosassociation of Colatoio Chemini and Laste Alte

This is the most havily trampled area of the whole ichnosite and is dedicated to the memory of Luciano Chemini, who discovered in 1989 the first dinosaur footprints of the site (Fig. 35).

The ichnoassemblage is very rich and comprises predominantly tridactyl footprints of small- to medium- sized theropod dinosaurs that fall in the ichnogenera *Kayentapus* isp. Welles, 1971, *Grallator* isp. Hitchcock, 1858 and *Eubrontes* isp. Hitchcock, 1845. *Kayentapus* isp. is the most common

morph. Some of the tridactyl footprints assigned to the ichnogenus *Anomoepus* isp. Hitchcock, 1848 have been attributed to basal ornithischians that had functionally tridactyl and clawed, theropod-like feet (Avanzini et al., 2001) but the interpretation remains uncertain. Some of the tridactyl tracks are elongate, displaying the whole or partial impressions of the metatarsus indicating a crouching posture of the dinosaur (Avanzini et al., 2001). In addition to tridactyl tracks there are many narrow-gauge trackways of medium-sized quadrupeds that document the presence of sauropodomorphs. Most of these trackways closely resemble those of *Parabrontopodus* Lockley, Farlow et Meyer, 1994 from the Upper Jurassic of Colorado (USA) and *Breviparopus* Dutuit et Ouazzou, 1980 from the Upper Jurassic–Lower Cretaceous of Morocco, both attributed to sauropodomorphs. Among the quadrupedal trackways, it is worth to mention the ichnotaxon *Lavinipes cheminii* Avanzini, Leonardi et Mietto, 2003 (Fig. 36) which was erected based on the unique morphologies shown by some large-sized tracks. *L. cheminii* has been attributed to basal sauropods (Avanzini et al., 2003).



Fig. 35: Several sauropod trackways can be seen along Colatoio Chemini.




Fig. 36: *Lavinipes cheminii* Avanzini, Leonardi et Mietto, 2003 (from Avanzini et al., 2003).

#### Stop 5, The Rotzo Formation and its exceptionally preserved flora

The Rotzo formation is characterised by a heterogeneous lithological composition (ooidal, peloidal, bioclastic and intraclastic limestones, marls, and clays), arranged into asymmetrical thickening and shallowing-upward meter-scale sequences (Masetti et al., 1998). This formation has been generally referred to a subtidal environment and can be subdivided into two main lithofacies: the lower is mainly represented by limestone-marl alternations, whereas the upper is characterized by the so-called *Lithiotis* beds or mounds (Masetti et al., 1998; Posenato & Avanzini, 2006). The lower lithofacies is essentially characterized by bioturbate mudstones-wackestones which are locally interbedded with microlaminated wackestones-packstones dark gray in color, slightly marly, strongly fissile, and with a high concentration of organic matter (Masetti et al., 1998; Bassi et al., 2008).

These black shale deposits are characterized by a well-preserved paucispecific and oligohaline fauna of small thin-shelled bivalves (*Eomiodon*, *Myrene*, and rare isognomids; Bassi et al., 2005), ostracodes (*Phraterfabanella*, *Klieana*, *Limnocythere*; Boomer et al., 2001), as well as fossil testate amoebae (*Diffugia*, *Pontigulasia*, *Centropyxis*; Bassi et al., 2008). Dinosaur tracks as well as vertebrate remains (crocodiloform teeth and semi-otid fish scales) have also been documented from these levels (Petti et al., 2011a, 2011b, 2013; M. Bernardi, F. M. Petti, and M. Avanzini, unpublished data). In addition, these black shales contain abundant macrofloral remains (cuticle, charcoal) and rhizoliths in life position, suggesting the presence of a well-developed flora, including woody plants (Avanzini, 1998; Masetti et al., 1998). The palynological assemblage is dominated by Circumpollen (56%), usually attributed to a xerophytic vegetation of subtropical, warm, and rather arid climate. *Azonotriletes* (40%; ferns) indicate

warm swamp or marsh vegetation, typical of a humid climate and freshwater influence. *Chasmatosporites* (2%; Cycadales) is typical of subtropical zones with warm-humid climates (see also Avanzini et al., 2006).

The flora of Rotzo is the most famous Jurassic flora of Italy. It has been studied since the XVIII century (e.g., Del Pozzo, 1764; Massalongo, 1851, 1853a, b, 1856, 1859). It was studied in detail by Barone Achille De Zigno (e.g., 1850, 1856–68, 1873–85, 1878) who described a total of 326 species (82 new species). Grandori (1913a, b, 1915) and Wesley (1956, 1958, 1966, 1974) reviewed the flora and identified several taxa of horsetails, ferns, seed ferns, cycadophytes and conifers. Horsetails are represented by *Equisetites* and *Phyllothea*. The ferns are represented by *Gleichenites*, *Dictyophyllum*, *Protorhipis*, *Hymenophyllites*,

*Coniopteris*, *Phlebopteris*, *Matonidium*, seed ferns by *Sagenopteris*, *Pseudosagenopteris*, *Cycadopteris* and *Dichopteris* while ginkgophytes are absent. *Sphenozamites*, *Otozamites*, *Zamites*, *Pterophyllum*, *Ptilophyllum*, *Weltrichia*, *Wielandiella* and *Blastolepis* are the cycadophyte genera in the flora. The conifers are represented by the genera *Brachyphyllum*, *Pagiophyllum*, *Elatocladus*, *Desmiophyllum*, *Pityophyllum* and *Dactylethrophyllum*. The flora is dominated by bennettitaleans and small-leaved conifers (e.g., Wesley, 1974; Barale, 1982; Kustatscher et al., 2014b) (Fig. 36). The reconstructed habitat corresponds to a brackish-swampy environment characterised by plants with coriaceous leaves, while conifers occupied the hinterland. The ferns also grew in areas that were not influenced by the brackish water. The climate was interpreted as semi-arid (Wesley, 1966).



Fig. 37: Some constituents of the Jurassic Rotzo flora. a) *Phyllothea* sp.; b) *Dichopteris* sp.; c) *Sphenozamites roessertii* (from Kustatscher et al., 2014b).

## References

- Abel, O. (1929): Eine Reptilien Fährte aus dem Oberen Perm (Grödener Sandstein) der Gegend von Bozen in Südtirol. – *Paleobiol.*, 2: 1–14, Wien.
- Allen, J.R.L. (1963): The classification of cross-stratified units, with notes on their origin. – *Sedimentology*, 2: 93–114.
- Avanzini, M. (1998): Resti di vertebrati dal Giurassico inferiore della Piattaforma di Trento (Italia settentrionale). Nota preliminare. – *Stud. Trent. Sci. Nat., Acta Geol.*, 73(1996): 75–80.
- Avanzini, M. (2000): *Synaptichnium* tracks with skin impressions from the Anisian (Middle Triassic) of the Southern Alps (Val di Non – Italy). – *Ichnos*, 7: 243–251.
- Avanzini, M. (2002a): The dinosaur site at the Lavini di Marco and the Lower Jurassic Calcarei Grigi Megatracksite. – In: Santantonio M. (ed.), *General Field Trip Guidebook, 6th International Symposium on the Jurassic System, 12–22.9.2002, Palermo, Italy*: 279–281.
- Avanzini, M. (2002b): Dinosauromorph tracks from the middle Anisian (middle Triassic) of the Southern Alps (Valle di Non - Italy). – *Boll. Soc. Pal. Ital.*, 41: 37–40.
- Avanzini, M., Bargossi, G.M., Borsato, A., Castiglioni, G.B., Cucato, M., Morelli, C., Prosser, G., Sapelza, A. (2007): Note illustrative della Carta Geologica d'Italia alla scala 1:50.000, foglio 026 Appiano. Provincia Autonoma di Trento, Provincia Autonoma di Bolzano. System Cart, Roma, 184 pp.
- Avanzini, M., Bargossi G.M., Borsato, A., Cucato, M., Morelli, C., Picotti, V., Selli L. (2012): Note illustrative della Carta Geologica d'Italia alla scala 1:50.000, foglio 043 Mezzolombardo. Servizio Geologico d'Italia – ISPRA, 256 p., Land Technology & Services, Roma.
- Avanzini, M., Bargossi, G.M., Borsato, A., Selli, L., (2010): Note Illustrative – Foglio 060 Trento, Carta Geologica d'Italia alla Scala 1:50000, p. 244, Roma, ISPRA.
- Avanzini, M., Bargossi, G.M., Castiglioni, G.B., Dalmeri, G., Eccel, E., Mancabelli, A., Morelli, C., Neri, C., Picotti, V., Prosser, G., Sartori, G., Zambotti, G. (2001): Carta Geologica della Provincia di Trento, tav. 26 III Fondo (a scala 1:25.000) con Note illustrative, 159 pp. Provincia Autonoma di Trento, Servizio Geologico.
- Avanzini, M., Bargossi, G.M., Cucato, M., Morelli, C., Prosser, G., Tomasoni, R. (2013): Foglio 026 Appiano. Guida ai percorsi geologici. Ufficio Geologia e prove materiali. Provincia Autonoma di Bolzano – Alto Adige, 324 pp., Litografia Artistica Cartografica – Firenze.
- Avanzini, M., Bernardi M., Nicosia, U. (2011): The Permo-Triassic tetrapod faunal diversity in the Italian Southern Alps. – In: Ahmad Dar, I., Ahmad Dar, M. (eds.): *Earth and Environmental Sciences*, 591–608, InTech.
- Avanzini, M., Frisia, S., Van den Driessche, K., Keppens, E. (1997): A Dinosaur Tracksite in an Early Liassic Tidal Flat in Northern Italy – Palaeoenvironmental Reconstruction from Sedimentology and Geochemistry. – *Palaios*, 12: 538–551.
- Avanzini, M., Gierlinski, G., Leonardi, G. (2001): First report of sitting *Anomoepus* tracks in European Lower Jurassic (Lavini di Marco site - Northern Italy). – *Riv. Ital. Pal. Strat.*, 107: 131–136
- Avanzini, M., Leonardi, G. (2002): *Isochirotherium inferi* ichnosp. n. in the Illyrian (Late Anisian, Middle Triassic) of Adige Valley (Bolzano – Italy). – *Boll. Soc. Paleont. Ital.*, 41: 41–50.
- Avanzini, M., Leonardi, G., Mietto, P. (2003): *Lavinipes cheminii* ichnogen., ichnosp. nov., a possible sauropodomorph tracks from the Lower Jurassic of the Italian Alps. – *Ichnos*, 10: 179–193.
- Avanzini, M., Lockley, M. (2002): Middle Triassic archosaur ontogeny and population structure: interpretation based on *Isochirotherium delicatum* fossil footprints (Southern Alps – Italy). – *Palaeogeo., Palaeoclim., Palaeoecol.*, 185: 391–402.
- Avanzini, M., Masetti, D., Romano, R., Podda, F., Ponton, M. (2007): Calcarei Grigi. – In: Cita Sironi, M.B., Abbate, E., Balini, M., Conti, M.A., Falorni, P., Germani, D., Groppelli, G., Manetti, P., Petti, F.M. (ed.), *Carta Geologica d'Italia – 1:50.000, Catalogo delle Formazioni, Unità tradizionali. Quaderni serie III, 7, Fascicolo VII. APAT, Dipartimento Difesa del Suolo, Servizio Geologico d'Italia*: 125–135.
- Avanzini, M., Mietto, P. (2008a): Lower and Middle Triassic footprints-based biochronology in the Italian Southern Alps. – *Oryctos*, 8: 3–13.
- Avanzini, M., Mietto, P. (2008b): The occurrence of the vertebrate ichnogenus *Synaptichium* in the Anisian (Middle Triassic) of Southern Alps. – *Studi Trent. Sci. Nat. Acta Geol.*, 83: 13–19.
- Avanzini, M., Neri, C. (1998): Impronte di tetrapodi da sedimenti anisici della Valle di Non (Trentino occidentale - Italia): nota preliminare. – *Ann. Mus. Civ. S. Nat. Ferrara*, 1: 5–19.

- Avanzini, M., Petti, F.M. (2008): Updating the dinosaur track-sites from the Lower Jurassic Calcarei Grigi Group (Southern Alps, Northern Italy). – *Stud. Trent. Sci. Nat., Acta Geol.*, 83: 289–301.
- Avanzini, M., Petti, F.M., Bernardi, M., Tomasoni, R. (2010): Crocodile-like footprints from the Upper Triassic (Carnian) of the Italian Southern Alps. – *New Mexico Mus. Nat. Hist. Sci. Bull.* 51, 61–64.
- Avanzini, M., Piubelli, D., Mietto, P., Roghi, G., Romano, R., Masetti, D. (2006): Lower Jurassic (Hettangian-Sinemurian) dinosaur track megasites, Southern Alps, Northern Italy. – In: Harris, J.D., Lucas, S.G., Spielmann, J.A., Lockley, M.G., Milner, A.R.C., Kirkland, J.I. (eds.), *The Triassic-Jurassic Terrestrial Transition*. – *New Mexico Mus. Nat. Hist. Sci. Bull.*, 37: 207–216.
- Avanzini, M., Renesto, S. (2002): A review of *Rhynchosauroides tirolicus* Abel, 1926 ichnospecies (Middle Triassic: Anisian-Ladinian) and some inferences on *Rhynchosauroides* trackmaker. – *Riv. Ital. Pal. Strat.*, 108: 51–66.
- Avanzini, M., Tomasoni, R. (2004): *Giornate di Paleontologia 2004, IV edizione, Bolzano 21-23 maggio 2004, Guida all'escursione: la gola del Bletterbach*. – *Stud. Trent. Sci. Nat., Acta Geologica*, 79 (2002): 1–34.
- Bain, R.J., Foos, A.M. (1993): Carbonate microfibrils related to subaerial exposure and paleosol formation. – In: Rezak, R., Lavoie, D.L. eds., *Carbonate Microfibrils*, 19–27. Springer Verlag, New York.
- Barale, G. (1982). Le genre *Cycadopteris* Zigno au Jurassique dans l'Ouest Européen. – *Palaeontographica*, B183(1–3): 8–56.
- Bassi, D., Fugagnoli, A., Masetti, D., Posenato, R. (2005). Stop 5. The Eomiodon assemblage of non-marine black shales: – In: Masetti, D., Posenato, R., Bassi, D., Fugagnoli, A. (eds.), *The Rotzo Formation (Lower Jurassic) at the Valbona Pass (Vicenza Province)*. – *Stud. Trent. Sci. Nat., Acta Geol., suppl. 1*, 80(2003): 25–27.
- Bassi, D., Fugagnoli, A., Posenato, R., Scott, D.B. (2008): Testate Amoebae from the Early Jurassic of the Western Tethys, North-East Italy. – *Palaeontology*, 51(6): 1335–1339.
- Bauer, K., Kustatscher, E., Butzmann, R., Fischer, T.C., Van Konijnenburg-van Cittert, J.H.A., Krings, M. (2014): Ginkgophytes from the upper Permian of the Bletterbach gorge (northern Italy). – *Riv. Ital. Paleont. Stratigr.* 120(3): 271–279.
- Bechstädt, T., Brandner R. (1971): Das Anis zwischen St. Vigil und dem Höhlensteintal (Pragser und Orlanger Dolomiten, Südtirol). – *Festband Geologisches Institut 300-Jahr-Feier Universität Innsbruck*: 9–103, Innsbruck.
- Benton, M.J., (2006): The origin of the dinosaurs. – *III Jornadas Internacionales sobre Paleontología de Dinosaurios y su Entorno, Colectivo Arqueológico-Paleontológico de Salas, Burgos, Spain*, p. 11–19.
- Benton, M.J., Twitchett, R.J. (2003): How to kill (almost) all life: the end-Permian extinction event. – *TREE*, 18: 358–365.
- Bernardi, M., Klein, H., Petti, F.M., Ezcurra, M.D. (2015): The origin and early radiation of archosauriforms: integrating the skeletal and footprint record. – *PLoS ONE* 10(6): e0128449.
- Bernardi, M., Kustatscher, E., Bauer, K., Petti, F.M., Franz, M., Wappler, T., van Konijnenburg-van Cittert, J.H.A. (2014): Reconstruction of a Loapingian (Late Permian) ecosystem from the Dolomites (Bletterbach, N Italy). – *Giornate di Paleontologia SPI 2014*, p. 23–24.
- Bernardi, M., Petti, F.M., Avanzini, M. (2010): A webbed archosaur footprint from the Upper Triassic (Carnian) of the Italian Southern Alps. – *New Mexico Mus. Nat. Hist. Sci. Bull.*, 51: 65–68.
- Bernardi, M., Petti, F.M., D'Orazi Porchetti, S., Avanzini, M. (2013): Large tridactyl footprints associated with a diverse ichnofauna from the Carnian of Southern Alps. – *New Mexico Mus. Nat. Hist. Sci. Bull.*, 61: 48–54.
- Bittner, A. (1892): Was ist norisch? – *Jb. k. k. Reichsanst.*, 42: 387–396.
- Boomer, I., Whatley, R., Bassi, D., Fugagnoli, A., Loriga, C. (2001): An early oligohaline ostracod assemblage within the marine carbonate platform sequence of the Venetian prealps, NE Italy. – *Palaeogeogr. Palaeoclimat. Palaeoecol.*, 166: 331–344.
- Borrouilh-Le Jean, F. (1993): Naissance Géologique d'un Karst. Lithification-dissolution des sédiments marins de 125 Ka à l'Actuel et ses conséquences (Gran Banc de Bahama et Tuamotu). – In: Maire R., Salomon J-N. (eds.), *Karst et évolutions climatiques*, 77–103. Université de Bordeaux, Bordeaux.
- Bosellini, A., Broglio Loriga, C. (1971): I «Calcarei Grigi» di Rotzo (Giurassico inferiore, Altopiano di Asiago) e loro inquadramento nella paleo-geografia e nella evoluzione tettonico-sedimentaria delle Prealpi venete. – *Ann. Univ. Ferrara, (N.S.)*, s. 9, 5/l: 1–61.
- Bosellini, A. (1984): Progradation geometries of carbonate platforms: examples from the Triassic of Dolomites, Northern Italy. – *Sedimentology*, 31: 1–24.
- Bosellini, A., Hardie, L.A. (1988): Facies e cicli della Dolomia Principale delle Alpi Venete. – *Mem. Soc. Geol. It.*, 30: 245–266.

- Brack, P., Kustatscher, E. (2013): *Voltzia recubariensis* from the uppermost Angolo Limestone of the Bagolino succession (Southern Alps of Eastern Lombardy, Italy). *Geo. Alp* 10: 61–70.
- Brack, P., Rieber, H., Nicora, A., Mundil, R., (2005): The Global boundary Stratotype Section and Point (GSSP) of the Ladinian Stage (Middle Triassic) at Bagolino (Southern Alps, Northern Italy) and its implications for the Triassic time scale. – *Episodes*, 28: 233–244.
- Breda, A. Preto, N. (2011): Anatomy of an Upper Triassic continental to marginal-marine system: the mixed siliciclastic-carbonate Travenanzes Formation (Dolomites, Northern Italy). – *Sedimentology*, 58: 1613–1647.
- Broglio-Loriga, C., Fugagnoli, A., Van Konijnenburg-van Cittert, J.H.A., Kustatscher, E., Posenato, R., Wachtler, M. (2002): The Anisian Macroflora from the Northern Dolomites (Kühwiesenkopf/Monte Pra della Vacca, Braies): a first report. – *Riv. It. Paleont. Strat.*, 108(3): 381–389.
- Brusatte, S.L., Nesbitt, S.J., Irmis, R.B., Butler, R.J., Benton, M.J. and Norell, M.A. (2010): The origin and early radiation of dinosaurs. – *Earth- Science Reviews*, 101: 68–100.
- Cassinis, G., Cortesogno, L., Gaggero, L., Massari, F., Neri, C., Nicosia, U., Pittau, P. (1999): Stratigraphy and facies of the Permian deposits between Eastern Lombardy and the Western Dolomites. – *Field Trip Guidebook, International Field Conference of "The Continental Permian of the Southern Alps and Sardinia (Italy). Regional Reports and General Correlations"*, pp. 157.
- Cassinis, G., Neri, C., Perotti, C.R. (1993): The Permian and the Permian-Triassic boundary in eastern Lombardy and western Trentino (Southern Alps, Italy). – *New Mexico Mus. Nat. Hist. Sci. Bull.* 3: 51–63.
- Castellarin, A., Piccioni, S., Prosser, G., Sanguinetti, E., Sartori, R., Selli, L. (1993): Mesozoic continental rifting and Neogene inversion along the South Giudicarie Line (Northwestern Brenta Dolomites). – *Mem. Soc. Geol. It.*, 49: 125–144.
- Castellarin, A., Picotti, V., Cantelli, L., Claps, M., Trombetta, L., Selli, L., Carton, A., Borsato, A., Daminato, F., Nardin, M., Santuliana, E., Veronese, L., Bollettinari, G. (2005): Note Illustrative della Carta Geologica d'Italia alla scala 1:50.000, foglio 080 Riva del Garda. – APAT, Dipartimento Difesa del Suolo, Servizio Geologico d'Italia, 145 pp.
- Ceoloni, P., Conti, M.A., Mariotti, N., Nicosia, U. (1988): New Late Permian tetrapod footprints from the Southern Alps. – In: Cassinis, G. (ed.), *Field Conference on Permian and Permian-Triassic boundary in the southalpine segment of the western Tethys, and additional regional reports*, Field conference, S.G.I., Brescia 4–12.07.1986, *Mem. Soc. Geol. It.*, 34(1986): 45–56.
- Clement-Westerhof, J.A. (1984): Aspects of Permian palaeobotany and palynology. IV. The conifer *Ortiseia* Florin from the Val Gardena Formation of the Dolomites and the Vicentinian Alps (Italy) with special reference to a revised concept of the Walchiaceae (Göppert) Schimper. – *Rev. Palaeobot, Palynol.*, 41: 51–166.
- Clement-Westerhof, J.A. (1986): Two peculiar types of coniferous ovuliferous fructifications from the Val Gardena Formation of the Dolomites and the Vicentinian Alps – a preliminary account. – *Cour. Forsch.-Inst. Senckenberg*, 86: 89–100.
- Clement-Westerhof, J.A. (1987): Aspects of Permian palaeobotany and palynology. VII The Majoniaceae, a new family of late Permian conifers. – *Rev. Palaeobot, Palynol.*, 52: 357–402.
- Conti, M.A., Leonardi, G., Mietto, P., Nicosia, U. (2000): Orme di tetrapodi non dinosauriani del paleozoico e Mesozoico d'Italia. – In: Leonardi G, Mietto P, editors. *Dinosauri in Italia*, Accademia Editoriale Pisa– Roma. pp. 297–320.
- Conti, M.A., Fontana D., Mariotti N., Massari F., Neri C., Nicosia U., Pasini M., Pittau P. (1986): The Bletterbach-Butterloch section (VGS and Bellerophon Formation). – In: *Italian IGCP Group 302 (ed.): Field Conference on Permian and Permian-Triassic boundary in the south-Alpine segment of the western Tethys*, Field guide book. S.G.I., 91–110, Brescia.
- Conti, M.A., Leonardi, G., Mariotti, N., Nicosia, U. (1975): Tetrapod footprints, fishes and molluscs from the Middle Permian of the Dolomites (N. Italy). – *Mem. Geopal. Univ. Ferrara*, 3(2): 139–150.
- Conti, M.A., Leonardi, G., Mariotti, N., Nicosia, U. (1977): Tetrapod footprints of the "Val Gardena Sandstone" (North Italy). Their paleontological, stratigraphic and paleoenvironmental meaning. – *Palaeont. It., N.S.*, 40: 1–91.
- Conti, M.A., Leonardi, G., Mariotti, N., Nicosia, U. (1980): Tetrapodi e pesci della Val Gardena. – In: AA.vv. (eds.): *I vertebrati fossili italiani - Catalogo della Mostra*, pp. 51–60, Verona.
- Cuny, G. Galton, P.M. (1993): Revision of the Airel theropod dinosaur from the Triassic-Jurassic boundary (Normandy, France). – *N. Jahrb. Geol. Paläont. Ab.*, 187: 261–288.
- D'Orazi Porchetti, S., Nicosia, U., Mietto, P., Petti, F.M. and Avanzini, M. (2008): *Atreipus*-like footprints and their co-occurrence with *Evazoum* from the upper Carnian (Tuvalian) of Trentino-Alto Adige. – *Studi Trentini Sci. Nat., Acta Geol.*, 83: 277–287.
- De Zanche, V., Franzin, A., Gianolla, P., Mietto, P., Siorpaes, C. (1992): The Piz da Peres section (Valdaora-Olang, Pusteria Valley). A reappraisal of the Anisian stratigraphy in the Dolomites. – *Eclog. Geol. Helv.*, 85: 127–143.

- De Zanche, V., Gianolla, P., Mietto, P., Siorpaes, C., Vail, P.R. (1993): Triassic sequence stratigraphy in the Dolomites (Italy). – *Mem. Sc. Geol.*, 45: 1–27.
- De Zigno, A. (1850): Coup d'oeil sur les Terrains stratifiés des Alpes Venetiennes. – 16 pp., W. Braumüller, Wien.
- De Zigno, A. (1856–68): Flora fossilis formationis oolithicae. Vol. 1. – 225 pp., Tip. Del Seminario, Padova.
- De Zigno, A. (1873–1885): Flora fossilis formationis oolithicae. Vol. 2. – 203 pp., Tip. Del Seminario, Padova.
- De Zigno, A. (1878): Sulla distribuzione geologica e geografica delle conifere fossili: memoria letta all'Accademia di scienze, lettere ed arti in Padova nella tornata del 7 luglio 1878. – 14 pp., Tipografia G.B. Brandi, Padova.
- Del Pozzo, A. (1764): Memorie storiche dei Sette Comuni vicentini. Vicenza
- Ezcurra, M.D., Scheyer, T.M., Butler, R.J. (2014): The Origin and Early Evolution of Sauria: Reassessing the Permian Saurian Fossil Record and the Timing of the Crocodile-Lizard Divergence. – *PLoS ONE*, 9(2): e89165.
- Farabegoli, E., Perri, M.C., Posenato, R. (2007): Environmental and biotic changes across the Permian–Triassic boundary in Western Tethys: the Bulla parastratotype, Italy. – In: Yin, H., Warrington, G., Xie, S. (eds.). *Environmental and Biotic Changes during the Paleozoic–Mesozoic Transition. Global and Planetary Change*, 55(1–3): 109–135.
- Fisher, J.A., Nichols, G.J., Waltham, D.A. (2007): Unconfined flow deposits in distal sectors of fluvial distributary systems: examples from the Miocene Luna and Huesca Systems, northern Spain. – *Sed. Geol.*, 195: 55–73.
- Gennaro M. (2007): The “Raibl Group” alongside the Adige Valley. – Università di Padova. Tesi di Laurea ined., A.A. 2006/2007: 144 p.
- Gianolla, P., Avanzini, M., Breda, A., Kustatscher, E., Preto, N., Roghi, G., Furin, S., Massari, F., Picotti, V., Stefani, M. (2010): DOLOMITES – 7th International Triassic Field Workshop (Pan–European Correlation of the Triassic). – Field trip to the UNESCO World Heritage Site of the Tethyan Triassic, 5–10.09.2010, Field trip guide book, 122 pp., Dolomites, Southern Alps, Italy.
- Gianolla, P., De Zanche, V., Mietto, P. (1998a): Triassic sequence stratigraphy in the Southern Alps (Northern Italy): definition of sequences and basin evolution. – In: de Graciansky P.C., Hardenbol J., Jacquin T., Vail P.R. (eds.): *Mesozoic–Cenozoic Sequence Stratigraphy of European Basins*, SEPM Special Publication 60: 723–751.
- Gianolla, P., Micheletti, C., Panizza, M., Viola, F. (2009): Nomination of the Dolomites for Inscription on the World Natural Heritage List Unesco: <http://whc.unesco.org/en/list/1237/>.
- Gianolla, P., Roghi, G., Ragazzi, E. (1998b): Upper Triassic amber in the Dolomites (Northern Italy). A paleoclimatic indicator? – *Riv. Ital. Paleont. Strat.*, 104: 381–390.
- Grandori, L. (1913a): Intorno alla flora dei Calcari grigi studiata da Achille De Zigno. – *Acc. Veneto-Trentino-Istria*, 14: 144–148.
- Grandori, L. (1913b): La Flora dei Calcari Grigi del Veneto. – *Mem. Ist. Geol. Univ. Padova*, 2: 1–112.
- Grandori, L. (1915): Su di un seme mesozoico di pteridosperma e sulle sue affinità con forme paleozoiche e forme viventi. – *Atti Acc. Veneto-Trentino-Istria*, 8: 107–116.
- Kerp, H., Kustatscher, E. (2005): The macroflora. – In: Pittau, P., Kerp, H., Kustatscher, E. (Eds.), *The Bletterbach Canyon: „Let us meet across the P/T boundary“*, Workshop on Permian and Triassic Palaeobotany and Palynology, 16–18.06.2005, Excursion guide, pp. 22–26, Bozen/Bolzano, Italy.
- Kittl, E. (1891): Saurier Färthe von Bozen. – *Mitt. Österreich. Tour. Klub*, 3: 7.
- Kreis, H.H. (1971): Sedimentologische Untersuchungen des „Unteren Muschelkalks“ (Anis) im Bereich des Gampen-Pass (Passo delle Palade) (Süd Tirol). – *Festb. Geol. Inst. Univ. Innsbruck*, 139–164.
- Kustatscher, E., Bauer, K., Butzmann, R., Fischer, T.C., Meller, B., Van Konijnenburg-van Cittert, J.H.A., Kerp, H. (2014a): Sphenophytes, seed ferns and possible cycads from the Wuchiapingian (Lopingian, Permian) of Bletterbach (Dolomites, Northern Italy). – *Rev. Palaeobot. Palynol.*, 208: 65–82.
- Kustatscher, E., Bauer, K., Reich, M. (2012): A new Middle Triassic (Pelsonian) plant locality in Northern Italy: The Non Valley (Trentino). – *Geo.Alp.* 9: 60–73.
- Kustatscher, E., Giusberti, L., Roghi, G. (2014b): La flora del Giurassico dell'Italia Settentrionale. – In: Kustatscher, E., Roghi, G., Bertini, A., Miola, A. (eds) *La storia delle piante fossili in Italia/Paleobotany of Italy*, pp. 154–165, Pubblicazione del Museo di Scienze Naturali dell'Alto Adige n. 9.
- Kustatscher, E., Van Konijnenburg-van Cittert, J.H.A., Bauer, K., Butzmann, R., Meller, B., Fischer, T.C. (2012): A new flora from the Upper Permian of Bletterbach (Dolomites, N-Italy). – *Rev. Palaeobot. Palynol.*, 182: 1–13.
- Land, L.S. (1986): Limestone diagenesis - Some geochemical considerations. – In: Mumpton, F.A. Ed., *Studies in Diagenesis*, U.S. Geol. Surv. Bul. 1578: 129–137.
- Leonardi, G., Lanzinger, M. (1992): Dinosauri nel Trentino: Venticinque piste fossili nel Liassico di Rovereto (Trento, Italia). – *Paleocronache*. 1: 13–24.
- Lee, M.S.Y. (1997): Pareiasaur phylogeny and the origin of turtles. – *Zool. J. Linn. Soc.*, 120: 197–280.

- Leonardi, G., Avanzini, M. (1994): Dinosauri in Italia. – Le Scienze, Quaderni, 76: 69–81.
- Leonardi, G., Mietto, P. (2000): Dinosauri in Italia: le orme giurassiche dei Lavini di Marco (Trentino) e gli altri resti fossili italiani. – Accademia editoriale. Pisa, Roma
- Leonardi, G., Nicosia, U. (1973): Stegocephaloid footprint in the Middle Permian sandstone (Groedener Sandsteine) of the Western Dolomites. – Ann. Univ. Ferrara, N.S., Sez. 9, 5(6): 116–1249.
- Leonardi, P. (1948): Contributi alla conoscenza della flora delle Arenarie di Val Gardena (Permiano medio-inf.) dell'Alto Adige: La nuova flora di Redagno e una felce di Egna. – Mem. Ist. Geol. Min. Univ. Padova, 16: 3–15.
- Leonardi, P. (1951): Orme di tetrapodi nelle Arenarie di Val Gardena (Permiano medio-inferiore) dell'Alto Adige sud-orientale. – Mem. Ist. Geol. Univ. Padova, 17: 1–23.
- Leonardi, P. (1953): La formazione desertica permiana delle Arenarie di Val Gardena e le loro flore e faune. – C.R. 19eme Sess. Congr. Géol. Intern., sez. 7: 33–36.
- Leonardi, P. (1968): Le Dolomiti. Geologia dei Monti fra Isarco e Piave. – 1019 pp., Manfrini Ed., Rovereto.
- Leonardi, P., Conti, M.A., Leonardi, G., Mariotti, N., Nicosia, U. (1975): *Pachypes dolomiticus* n.gen. n.sp.: Peirosaur footprint from the "Val Gardena Sandstone" (Middle Permian) in the western Dolomites (N. Italy). – Atti Acc. Naz. Lincei, Rend. Cl. Sci. ff. mm. nn., ser 8, 57: 221–232.
- Lockley, M. (2002): Fossil Footprints of the World. – 124 pp., University of Colorado, Denver.
- Lucas, S.L., Klein, H., Lockley, M.G., Spielmann, J.A., Gierlinski, G.D., Hunt, A.P., Tanner, L.H., (2006): Triassic-Jurassic stratigraphic distribution of the theropod footprint ichnogenus *Eubrontes*. – New Mexico Mus. Nat. Hist. Sci. Bull., 37: 86–93.
- Masetti, D., Bottoni, A. (1978): L'Encrinite di Fanes e suo inquadramento nella paleogeografia giurassica dell'area Dolomitica. – Riv. Ital. Paleont., 84: 169–186.
- Masetti, D., Claps, M., Giacometti, A., Lodi, P., Pignatti, P., (1998): I Calcari Grigi della Piattaforma di Trento (Lias Inferiore e Medio, Prealpi Venete). – Atti Tic. Sci. Terra, 40: 139–183.
- Massalongo A. (1853a): *Plantae fossiles novae in formationibus tertiaris regni veneti nuper inventae*. – Tip. Ramanzini, Verona.
- Massalongo A. (1856): Analisi di una lignite scavata nel comune di Badia Calavena nel veronese. – Estratto dalla Gazz. Uff. di Verona, Anno II, N. 77.
- Massalongo, A. (1851): Sopra le piante fossili dei terreni terziari del vicentino. – Tipografia A. Bianchi, Padova.
- Massalongo, A. (1853b): Enumerazione delle piante fossili Miocene fino ad ora conosciute in Italia. – 31 pp., G. Antonelli, Verona.
- Massalongo, A. (1859): Specimen photographicum animalium quorundam plantarumque fossilium Agri Veronensis. A. B. Massalongo descripsit Mauritius Lotze photographice expressit. – 101 pp., Typ. Vicentini Franchini, Verona.
- Massari, F., Neri, C. (1997): The infill of a supradetachment (?) basin: the continental to shallow-marine Upper Permian in the Dolomites and Carnia (Italy). – Sed. Geol., 110: 181–221.
- Massari, F., Neri, C., Fontana, D., Manni, R., Mariotti, N., Nicosia, U., Pittau, P., Spezzamonte, M., Stefani, C. (1999): The Bletterbach section (Val Gardena Sandstone and Bellerophon Formation). – In: Cassinis, G., Cortesogno, L., Gaggero, L., Massari, F., Neri, C., Nicosia, U., Pittau, P. (eds.), Stratigraphy and facies of the Permian deposits between eastern Lombardy and the Western Dolomites. Field Trip Guidebook, pp. 111–132, Earth Science Department, Pavia.
- McKenzie, J.A., Vasconcelos, C. (2009): Dolomite Mountains and the origin of the dolomite rock of which they mainly consist: historical developments and new perspectives. – Sedimentology, 56: 205–219.
- Miall, A.D. (1985): Architectural-element analysis: a new method of facies analysis applied to fluvial deposits. – Earth Science Review, 22: 261–308.
- Mietto, P., Gianolla, P., Manfrin, P., Preto, N. (2003): Refined ammonoid biostratigraphy of the Bagolino section (Lombardian Alps, Italy), GSSP candidate for the base of the Ladinian Stage. – Riv. Ital. Paleontol. S., 109(3): 449–462.
- Mietto, P., Manfrin, S. (1995): A high resolution Middle Triassic ammonoid standard scale in the Tethys Realm. A preliminary report. – Bull. Soc. Géol. France, 166: 539–563.
- Mietto, P., Manfrin, S., Preto, N., Rigo, M., Roghi, G., Furin, S., Gianolla, P., Posenato, R., Muttoni, G., Nicora, A., Buratti, N., Cirilli, S., Spötl, C., Ramezani, J., Bowring, S.A. (2012): The Global Boundary Stratotype Section and Point (GSSP) of the Carnian Stage (Late Triassic) at Prati di Stuores/Stuores Wiesen Section (Southern Alps, NE Italy). – Episodes, 35(3): 414–420.
- Mojsisovics, E. von (1879): Die Dolomit-Riffe von Südtirol und Venetien. Beiträge zur Bildungsgeschichte der Alpen. – 552 pp., Ed. Hödler, Wien.
- Mojsisovics, E. von (1882): Die Cephalopoden der mediterranen Triasprovinz. – Abh. k.-k. geol. Reichsanst., 10: 1–322.

- Mojsisovics, E. von, Waagen, W., Diener, C. (1895): Entwurf einer Gliederung der pelagischen Sedimente des Trias-Systems. – Sitzber. Akad. Wiss., math.-natw. Kl., 104(1): 1271–1302.
- Morelli, C., Bargossi, G.M., Mair, V., Marocchi, M., Moretti, A. (2007): The lower Permian volcanics along the Etsch valley from Meran to Auer (Bozen). – Mitt. Österr. Miner. Ges., 153: 195–218.
- Neri, C., Gianolla, P., Avanzini, M. (2005): Raibl beds (Upper Triassic): A case-history about the use of traditional lithostratigraphic names from central-eastern southern Alps. – *Geoitalia 2005*, V Forum Italiano di Scienze della Terra, 1: 9.
- Orombelli, G., Sauro, U. (1989): I Lavini di Marco: un gruppo di frane Oloceniche nel contesto morfotettonico dell'alta Val Lagarina (Trentino). – Suppl. Geog. Fis. Din. Quat., 1: 107–116.
- Perwanger, L. (1946): Geologisch Beobachtenswertes auf der Hochfläche von Radein. – *Der Schlern*, 20: 362–370.
- Petti, F.M., Bernardi, M., Todesco, R., Avanzini, M. (2011a): Vertebrate palaeontology and palaeoenvironment of the lower Rotzo Formation (Early Jurassic, Trento carbonate platform, NE Italy). – *Giornate di Paleontologia SPI 2011*, Abstract volume.
- Petti, F.M., Bernardi, M., Todesco, R., Avanzini, M. (2011b): Dinosaur footprints as ultimate evidence for a terrestrial environment in the Late Sinemurian Trento Carbonate Platform. – *Palaios*, 26: 601–606.
- Petti, F.M., Bernardi, M., Ferretti P., Tomasoni, R., Avanzini, M. (2011c): Dinosaur tracks in a marginal marine environment: the Coste dell'Anglone ichnosite (Early Jurassic, Trento Platform, NE Italy). – *Italian Journal of Geosciences*, 130 (1), 27–41 (doi: 10.3301/IJG.2010.19).
- Petti, F.M., Bernardi, M., Kustatscher, E., Renesto, S., Avanzini, M. (2013): Diversity of continental tetrapods and plants in the Triassic of the Southern Alps: ichnological, paleozoological and paleobotanical evidence. – *New Mexico Museum of Natural History and Science, Bulletin*, 61: 458–484.
- Pittau, P., Kerp, H., Kustatscher, E. (2005): The Bletterbach canyon. – “Let us meet across the P/T boundary” – workshop on Permian and Triassic Palaeobotany and Palynology, Bozen 16.–18.06.2005, Excursion guide, pp. 39, Bozen.
- Piubelli, D., Avanzini, M., Mietto, P. (2005): The Early Jurassic ichnogenus *Kayentapus* at Lavini di Marco ichnosite (NE Italy). Global distribution and palaeogeographic implication. – *Boll. Soc. Geol. It.*, 124: 259–267.
- Poort, R.J., Kerp, J.H.F. (1990): Aspects of Permian palaeobotany and palynology. XI. On the recognition of true peltasperms in the Upper Permian of western and central Europe and a reclassification of species formerly included in *Peltaspermum* Harris. – *Rev. Palaeobot. Palynol.*, 63: 196–225.
- Posenato, R. (2010): Marine biotic events in the Lopingian succession and latest Permian extinction in the Southern Alps (Italy). – *Geol. J.*, 45: 195–215.
- Posenato, R., Avanzini, M., (2006). “Lithiotis” beds of the Rotzo Formation (Calcarei Grigi Group, Lower Jurassic) from Albaredo (Rovereto, Trento): – *Stud. Trent. Sci. Nat., Acta Geol.*, 81(2004): 23–28.
- Posenato, R., Holmer, L.E., Prinoth, H. (2014). Adaptive strategies and environmental significance of lingulid brachiopods across the late Permian extinction. – *Palaeogeog. Palaeoclim. Palaeoecol.* 399: 373–394.
- Renesto, S., Bernardi, M. (2014): Redescription and phylogenetic relationships of *Megachirella wachtleri* Renesto et Posenato, 2003 (Reptilia, Diapsida). – *Paläont. Zeitschrift* 88: 197–210.
- Renesto, S., Posenato R. (2003): A new lepidosauromorph reptile from the Middle Triassic of the Dolomites. – *Riv. Ital. Pal. Strat.*, 109: 463–474.
- Resch, W. (1979): Zur Fazies-Abhängigkeit alpiner Trias-Foraminiferen. – *Jahrb. Geol. B.-A.*, 122(1): 181–249.
- Richthofen, F. (1860): Geognostische Beschreibung der Umgebung von Predazzo Sanct Cassian und Seiseralpen in Südtirol. – 327 pp., Ed. Perthes, Gotha.
- Roghi, G. (2004): Palynological investigation in the Carnian of the Cave del Predil area (Julian Alps, Italy). – *Rev. Palaeobot. Palynol.*, 73: 774–89.
- Roghi, G., Kustatscher, E., Van Konijnenburg-van Cittert, J.H.A. (2006): Late Triassic Plant from Julian Alps (Italy). – *Boll. Soc. Paleont. It.*, 45: 133–140.
- Roghi, G., Kustatscher, E., Bernardi, M., Dal Corso, J., Forte, G., Franz, M., Hochuli, P., Krainer, K., Petti, F.M., Ragazzi, E., Riva, A., Wappler, T., Gianolla, P. (2014): Field trip to Permo-Triassic palaeobotanical and palynological sites of the Southern Alps. – *Geo.Alp*, 11: 29–84.
- Sauro, U., Meneghel, M. (ed., 1995): Altopiani Ampezzani, geologia, geomorfologia e speleologia. – 156 pp., Ed. La Grafica, Cortina.
- Schmidt, A.R., Ragazzi, E., Coppellotti, O., Roghi, G. (2006): A microworld in 220 million-year-old drops of amber. – *Nature*, 444: 835.
- Valdiserri, D., Avanzini M. (2007): A Tetrapod Ichnoassociation from the Middle Triassic (Anisian, Pelsonian) of Northern Italy. – *Ichnos*, 14:105–116.

- Valentini, M., Conti, M.A., Nicosia, U. (2008): Linking tetrapod tracks to their trackmakers' bio- dynamics, paleo- biogeography, and paleobiology: *Pachypes dolomiticus* Leonardi et al., 1975, a case study. – Stud. Trent. Sci. Nat., Acta Geol., 83: 237–256.
- Valentini, M., Nicosia, U., Conti, M.A. (2009): A re-evaluation of *Pachypes*, a pareiasaurian track from the Late Permian. – N. Jb. Geol. Paläont. Abh., 251: 71–94.
- Venzo, G.A. (2000): Inquadramento geografico, geologico e cronistoria degli studi e ricerche sui Lavini di Marco. – In: Leonardi G., Mietto P. (eds.), Dinosauri in Italia. pp. 21–32, Accademia Ed., Pisa-Roma.
- Wesley, A. (1956): Contributions to the Knowledge of the flora of the Grey Limestones of Veneto, Part I. A revision of the Flora Fossilis formationis Oolithicae of De Zigno. – Mem. Ist. Geol. Min. Univ. Padova, 19: 1–69.
- Wesley, A. (1958): Contributions to the Knowledge of the flora of the Grey Limestones of Veneto, Part II. A revision of the Flora Fossilis formationis Oolithicae of De Zigno. – Mem. Ist. Geol. Min. Univ. Padova, 21: 1–57.
- Wesley, A. (1966): The fossil flora of the Grey Limestones of Veneto, Northern Italy, and its relationships to the other European floras of similar age. – The Palaeobotanist, 14: 124–130.
- Wesley, A. (1974): On the Bennettitalean remains from the Lias of Northern Italy. In: Symposium on morphological and stratigraphical Palaeobotany. – Birbal Sahni Institute of Palaeobotany Special Publication (Lucknow, Gumti) 2: 66–72.
- Wopfner, H. (1999): Über Tetrapoden-Fährten, Kohlen und versteinerte Holzer aus dem Grödner Sandstein (Perm) von Deutschnofen. – Der Schlern, 73: 23–32.
- Zenger, D.H., Bourrouilh-Le Jan, F.G., Carozzi, A.V. (1994): Dolomieu and first description of dolomite. – Spec. Publs. Int. Ass. Sedimentol., 21: 21–28.

